



**Verde Valley Medical Center**  
Northern Arizona Healthcare

## **Prehospital Care Treatment Guidelines**

---

**Carrie Burns M.D.**  
**VVMC Base Hospital Administrative Medical Director**  
**Revised April, 2011**

---

**Date**

<b><u>TABLE OF CONTENTS</u></b>	<b><u>PG</u></b>
<b>Introduction</b>	<b>7</b>
<b>Goals of Pre-Hospital Care</b>	<b>7</b>
<b>Medical Control</b>	<b>7</b>
<b>Standing Orders</b>	<b>8</b>
<b>Medical Control Options</b>	<b>8</b>
<b>Determination of Death</b>	<b>8</b>
<b>Withholding /Termination of Resuscitation</b>	<b>9</b>
<b>Healthcare Directives</b>	<b>9</b>
<b>Medical Control of A.L.S. at the Scene</b>	<b>9</b>
<b>Medical Cases</b>	<b>10</b>
<b>Trauma Cases</b>	<b>10</b>
<b>Obstetrical-Gynecological Cases</b>	<b>10</b>
<b>Psychiatric Cases</b>	<b>10</b>
<b>General Cases</b>	<b>11</b>
<b>Communications</b>	
<b>General Procedure</b>	<b>11</b>
<b>Definitions</b>	
<b>ALS Stable Situation</b>	<b>12</b>
<b>ALS Unstable Situations</b>	<b>12</b>
<b>Courtesy Notification</b>	<b>13</b>
<b>Exceptions</b>	<b>13</b>
<b>Basic Radio Procedures</b>	<b>14</b>
<b>Communication Guidelines</b>	<b>14</b>
<b>Communications Systems Failures</b>	<b>14</b>
<b>Intermediary's Responsibility</b>	<b>15</b>
<b>Body Substance Isolation</b>	<b>15</b>
<b>Transportation</b>	<b>15</b>

<b>Interfacility Patient Transportation</b>	<b>16</b>
<b>At Scene Transfer of Patient Care</b>	<b>16</b>
<b>Trapped or Impaled Patient</b>	<b>17</b>
<b>Refusal of Transport</b>	<b>17</b>
<b>Field Triage Guidelines</b>	<b>18</b>
<b>Multiple Casualty Incidents (M.C.I.)</b>	<b>18</b>
<b>Pneumatic Anti-Shock Garment (P.A.S.G.) Guidelines</b>	<b>20</b>
<b>Treatment Guidelines</b>	
<b>General Assessment and Treatment Approach</b>	<b>20</b>
<b>History</b>	<b>20</b>
<b>Initial Assessment</b>	<b>20</b>
<b>Airway</b>	<b>20</b>
<b>Breathing</b>	<b>21</b>
<b>Circulation</b>	<b>21</b>
<b>Vital Signs</b>	<b>21</b>
<b>Neurological Assessment</b>	<b>21</b>
<b>General: Focused History/Physical Exam or Rapid Assessment</b>	
<b>Detailed Physical Exams- Definitions</b>	<b>22</b>
<b>Central Venous Line Access- Paramedics Only</b>	<b>23</b>
<b>CPAP</b>	<b>23</b>
<b>Intravenous IV Access/IO</b>	<b>23</b>
<b>Rapid Sequence Intubation/RSI/ Medication Assisted Intubation</b>	<b>23</b>
<b>TREATMENT ALGORITHM INDEX</b>	
<b>DOA/DNR:</b>	
<b>Dead on Arrival</b>	<b>24</b>
<b>Do Not Attempt Resuscitation</b>	<b>25</b>
<b>Withholding/Termination of Resuscitation</b>	<b>26</b>
<b>CARDIAC:</b>	
<b>Adult Bradycardia, Unstable</b>	<b>27</b>
<b>Adult Chest Pain suggestive of cardiac origin</b>	<b>28</b>

Adult Cardiopulmonary Arrest- CCR Alternative	29
Adult Pulseless Arrest	30
Cardiac Arrest Post Resuscitation Induced Hypothermia	31
Adult Tachycardia with Pulses	32
<b>TRAUMA</b>	
Trauma Triage Designation	33
Submersion Incident	34
Trauma-Burns	35
Trauma-Musculoskeletal Injury	36
Trauma-Head Injury with ALOC	37
Trauma-Multi-System	38
Trauma-Spinal Injury	39
<b>CVA</b>	
Cerebral Vascular Accident-Stroke	40
<b>ALLERGIC REACTIONS</b>	
Allergic Reaction	41
Anaphylaxis	42
Envenomation-Arachnids	43
Envenomation-Snake Bites	44
<b>AIRWAY</b>	
Airway Compromise	45
Airway Obstructed	46
Respiratory Insufficiency-Bronchospasm	47
Respiratory Insufficiency-Pulmonary Edema	48
<b>OBSTETRICS</b>	
Complications of Delivery Abnormal Presentations	49
Complications of Delivery Post Partum Hemorrhage	50
Complications of Pregnancy	51
Complications of Pregnancy Cont	52
Delivery	53
<b>ALOC/SEIZURES</b>	
Altered Level of Consciousness	54
Altered Level of Consciousness with ETOH	55
Adult Violent/Agitated Patient	56
Seizure	57
<b>MEDICAL</b>	
Abdominal Pain, Non-Traumatic	58
Nausea and Vomiting	59

Environmental-Heat Related	60
Environmental-Hypothermia	61
Hypotension, Non-Traumatic	62
Poisoning/Overdose	63
<b>PEDIATRICS</b>	
Ped. Bradycardia, Unstable	64
Ped. Pulseless Arrest	65
Ped. Tachycardia with Pulses	66
Ped. Neonatal Resuscitation	67
Ped. Submersion Incident-Category 1	68
Ped. Submersion Incident-Category 2	69
Ped. Trauma-Burns	70
Ped. Trauma-Musculoskeletal Injury	71
Ped. Trauma-Head Injury with ALOC	72
Ped. Trauma-Multi-System	73
Ped. Trauma-Spinal Injury	74
Ped. Allergic Reaction	75
Ped. Anaphylaxis	76
Ped. Envenomation –Arachnids	77
Ped. Envenomation-Snake Bites	78
Ped. Airway Compromise	79
Ped. Airway Obstructed	80
Ped. Insufficiency-Bronchospasm	81
Ped. Respiratory-Upper Airway Emergencies Croup/Epiglottitis	82
Ped. ALOC	83
Ped. Seizures of Unknown Etiology	84
Ped Nausea and Vomiting	85
Ped. Abdominal Pain, Non-Traumatic	86
Ped. Environmental-Heat Related	87
Ped. Environmental-Hypothermia	88
Ped. Hypotension/Shock, Non-Traumatic	89
Ped. Poisoning/Overdose	90
<b>Appendix</b>	
Appendix A- Pediatric/Neonatal Standards/Pharmacological Modalities	91
Appendix B- Adult Pharmacological Modalities	94
Appendix C- Prehospital Standard Infusion Mixtures	96
Appendix D- Authorized Medication/Interfacility Transport List	97
Appendix E- Scores/Scales/Questionnaires	101
• Glasgow Coma Scale	
• Pediatric Glasgow Coma Scale	
• F.A.S.T Stroke Scale	
• APGAR Score	
Appendix F- Transport Guidelines	105

- **VVEMS Medical Direction Policy on Transport Destination**
- **Sedona Transportation Guidelines**

<b>Appendix G- RSI, CPAP, IO ACCESS</b>	<b>107</b>
<b>Appendix H- Head to Toe Assessment</b>	<b>113</b>
<b>Appendix I- ALS Release of Patients for BLS Transport</b>	<b>114</b>
<b>Appendix J – EMTI Care Levels</b>	<b>115</b>

## **INTRODUCTION**

The purpose of these treatment guidelines is to provide uniform prehospital care for agencies under the medical direction of Verde Valley Medical Center Base Hospital (referred to as VVEMS Agencies). They are directed towards A.L.S. (IEMT99/CEP) levels of Arizona Department of Health Services (A.D.H.S) certified pre-hospital care providers.

## **GOALS OF PRE-HOSPITAL CARE**

The first goal of pre-hospital care is on-scene recognition and treatment of conditions in which the delay of treatment might increase morbidity and mortality. Once the patient enters the Emergency Medical Services (EMS) system, life-saving interventions should be initiated immediately.

The second goal is rapid transport, with only minimal on-scene delay, for patients whose conditions require immediate hospital stabilization.

The third goal of pre-hospital care is to provide initial stabilization, safe symptom relief and safe transport to a medical facility.

The fourth goal is on-scene triage in multiple casualty incidents.

To achieve the above stated goals of pre-hospital care, the medic must be skilled in patient assessment. He/she must be able to recognize those conditions where on-scene intervention is necessary and those when rapid transport is best.

Assessment must be rapid, succinct and goal directed. Main emphasis is on the primary survey. Secondary survey should not delay either life saving interventions or transport. Interventions identified in the assessment should be acted on immediately.

## **MEDICAL CONTROL**

It is important to recognize that emergency care rendered in the pre-hospital environment, even though performed by an emergency medical technician, remains the responsibility of the On-line Physician. These treatment guidelines are not intended for use as inflexible rules for pre-hospital care, but rather as guidelines for physicians and pre-hospital care personnel alike. Although they represent a minimum standard of care against which actions may be judged, treatment guidelines are not absolute. Common sense and good judgment are equally important. Since individual situations may require variance from these guidelines, the final authority is the independent medical judgment of the medical control physician. Also, it should be understood that skill levels of individuals will vary, and the online medical control may find it necessary to vary from these guidelines. EMS providers are expected to use online medical control as a real time consultant when there are any doubts or concerns as to what is the correct course of action.

## **STANDING ORDERS**

Standing orders are those interventions, approved by the Administrative Medical Director, which may be done immediately, prior to radio contact with online medical control. Generally, they will include those life or limb saving procedures where either the delay caused by radio communication could contribute to death or where there is no disagreement about what should be done in a very specific situation or treatments with a consistently favorable risk/benefit ratio.

## **MEDICAL CONTROL OPTIONS**

Medical control option means that the procedure requires a specific order from the online medical control via radio or telephone prior to performance. Any situation where procedures are performed, which by these treatment guidelines require a medical control option, and such medical control option is not obtained because of inability to establish radio contact or due to the critical nature of the situation, clear cut indications for the procedure(s) must exist (according to the treatment guidelines herein). We do not wish patients to suffer because of inadequacies or failures of the communication system but patient safety is of great importance. Communication with the Base Hospital should be established as soon as possible in such incidents.

Medical control options will be noted as footnotes in the individual treatment guidelines.

## **DETERMINATION OF DEATH**

Prehospital personnel respond to victims of cardiopulmonary arrest in a variety of circumstances. The following guidelines are intended to assist in determining how and when resuscitative measures should be withheld, initiated, and/or terminated. Refer to appropriate related treatment algorithms for other specific information.

If the victim meets the criteria listed below, no resuscitative efforts need to be initiated. On-line medical control is NOT necessary. Contact law enforcement and initiate grief support. An EMS provider must remain with the victim until released to law enforcement.

All of the following criteria must be met:

- Patient is pulseless and apneic
- Presence of one or more signs of irreversible death
- Asystole is confirmed on the monitor in two leads for at least 12 seconds as defined in the guideline
- Hypothermia is not present

Signs of irreversible death:

- Decapitation
- Decomposition
- Transection of thorax or abdomen
- Burned beyond recognition
- Dependent Lividity and/or rigor mortis and Asystole in 2 leads for 12 seconds

## **WITHOLDING/TERMINATION OF RESUSCITATION**

Field termination of resuscitative efforts may be considered for both trauma and medical patients. Patients must be in cardiopulmonary arrest in a rhythm incompatible with life (asystole, pulseless electrical activity). Treat patients according to the trauma or medical field termination of guideline and associated treatment algorithm. On-line medical direction is required for all field terminations.

## **HEALTHCARE DIRECTIVES**

If a valid Prehospital Medical Care Directive is present, no resuscitative measures are needed. A patch should be done to the base hospital if possible.

If a valid Living Will/Advanced Directive/Do Not Resuscitate consent or orders is present, begin resuscitation and contact medical control.

## **MEDICAL CONTROL OF Advanced Life Support (A.L.S.) AT THE SCENE**

General Principles:

When an A.L.S. unit, under medical direction, is requested and dispatched to the scene of an emergency, a doctor/patient relationship has been established between the patient and the physician providing medical direction. The individual with the highest level of certification is responsible for management of the patient, and acts as the agent of medical direction unless the patient's physician is present.

If the patient's private physician is on the scene or a physician intervener\* is present and he/she prefers to assume responsibility for care, the On-line Physician must be contacted and the situation discussed. Only Medical Control can relinquish care of the patient to another physician. Any action performed by the medic at the physician intervener's direction must be in line with local treatment guidelines. If not, Medical Control should be contacted. In any event, the physician intervener is responsible for appropriate documentation and, unless absolute necessity dictates otherwise, should accompany the patient to the hospital.

Intervener physician is a licensed physician who has not established a prior physician/patient relationship and who wishes to take charge of a medical emergency scene, and who is willing to provide evidence of licensure and agrees to continue care for the patient during transport to the hospital if feasible.

If an intervener physician is present and on-line medical direction does exist, the On-line Physician is ultimately responsible. If there is any disagreement between the intervener physician and the on-line physician, medical direction will remain with medical control. The on-line physician has the option of managing the case entirely, working with the intervener physician, or allowing him to assume responsibility. In the event that the intervener physician assumes responsibility, all orders to the A.L.S. provider should be repeated over the radio for

purposes of recording. The intervener physician should document his intervention in a manner acceptable to the local E.M.S. system. The decision of the intervener physician to accompany the patient to the hospital should be made in consultation with the on-line physician. If on-line medical direction is not possible, treatment guidelines will be followed.

## **ALS CALLS**

A.L.S. providers shall contact the On-line Physician for medical direction, as defined in the treatment guidelines.

## **MEDICAL CASES**

- Chest Pain
- Shortness of breath
- Hematemesis, melena, or hematochezia
- Altered Level of Consciousness
- Loss of consciousness (syncope, seizures)
- Possible drug overdose or ingestion of poisonous substances
- Recent change in mental status
- More than one acutely ill person
- Painful, cold, pulseless, extremity
- Acute abdominal pain
- Terminal malignancy in distress

## **TRAUMA CASES**

- Motorcycle, auto vs. pedestrian or bicycle accidents
- Suspected fractures of femur, pelvis, spine, or skull
- Extremity wounds with distal neurological and/or vascular compromise
- Head injuries with history of loss of consciousness or presently impaired mental status
- Penetrating wounds of head, neck, chest, abdomen, or thigh
- Blunt trauma to abdomen or chest wall
- Burn Injuries
- Significant acute external blood loss
- Water accidents and near drowning
- Extrication problems
- Multiple casualties

## **OBSTETRICAL-GYNECOLOGICAL CASES**

- Vaginal Hemorrhage
- Childbirth
- Pregnancy with abdominal pain

## **PSYCHIATRIC CASES**

- Suicide (attempts or verbalization)
- Hallucinations with behavioral problem
- Violent or dangerous patients (result of mental disorders)

## **GENERAL CASES**

Signs of shock

Hypotension (systolic blood pressure of 90 or less in an adult) significant tachycardia or other abnormal vitals

Altered mental status

D.O.A. patients

Any patient who, in the opinion of the A.L.S. personnel, would benefit from Base Hospital consultation.

Any patients with suspected medical or traumatic problems of an A.L.S. nature, who refuse treatment or transportation to a hospital.

Abnormal body temperatures

When there is a physician on the scene who wishes to take control of patient care.

## **A.L.S. RESPONDERS MUST PATCH IN ANY QUESTIONABLE OR UNUSUAL SITUATION**

Environmental hazard

Security problem

When disagreements arise between responding E.M.S. providers or with law enforcement

## **COMMUNICATIONS**

### **GENERAL PROCEDURE:**

Participating A.L.S. providers shall initiate ALS care through the use of treatment guidelines, and dependent upon patient response or treatment guideline criteria shall have the following communication options:

1. Stable Situation:
  - a. Courtesy Notification (CN) with Receiving Facility
  - b. Courtesy Notification (CN) with Base Hospital (to have information relayed to receiving facility)
  - c. Patch with Base Hospital
2. Unstable Situation after implementation of standing orders:
  - a. Patch with Base Hospital
3. Exception Situations:
  - a. Critical Trauma, Medical Code:
    - i. Courtesy Notification (CN) with Receiving Facility
    - ii. Courtesy Notification (CN) with Base Hospital (to have information relayed to receiving facility)
    - iii. Patch with Base Hospital
  - b. Unable to contact Base Hospital:
    - i. Patch with designated back-up for Base Hospital
    - ii. Patch with Receiving Facility

## **DEFINITIONS:**

### **1. ALS STABLE SITUATION (Requires minimum of Courtesy Notification):**

All patients are assumed to be ALS unless criteria for BLS are present and the providers and online medical direction are comfortable making the patient a BLS transport. This will require a patch by medic requesting permission to down grade pt to BLS. SEE APPENDIX J for ALS Release of Patients for BLS Transport.

A patient with a single system or well-defined chief complaint(s) that after initial ALS intervention is:

- Without neurological, respiratory and/or cardiovascular compromise; or
- Has responded favorably to initial treatment modalities (resolving or improving chief complaint and/or signs/symptoms).

#### **Criteria for ALS Stable Situations may include:**

- a. Conscious, alert and oriented to person, time, place and event (with consideration of pre-existing conditions) or an altered mental status in a non-traumatic event after treatment with no signs of impending central herniation, GCS maintained at  $\geq 14$  and stable vital signs.
- b. Respirations within normal range for age group and without abnormal breath sounds (with consideration of preexisting conditions).
- c. Pulse within normal range for age group and without irregularities (with consideration of preexisting conditions).
- d. Blood pressure greater than 90 systolic and less than 180 systolic, or within normal range for age group (with consideration of pre-existing conditions).
- e. No uncontrolled bleeding.
- f. Relief of chest pain.

### **2. ALS UNSTABLE SITUATIONS (Requires Patch):**

A patient with a single or multiple system or complex chief complaint with/without hemodynamic compromise and that does not respond favorably to initial treatment modalities. Refer to “Exceptions” for Critical Trauma and Medical Codes. Criteria for an unstable patient condition may be indicated by the presence of any of the following:

- a. Any situation where management is uncertain or risk benefit ratio of intervention is unclear or provider feels that patient is unstable or may deteriorate en route.
- b. ALOC, adult or (with consideration of pre-existing conditions). pediatric all causes other than resolving postictal signs and symptoms (S/S).
- c. Abnormal blood pressure (with consideration of pre-existing conditions).
- d. Abnormal heart rate or rhythm persisting after treatment that is causing hemodynamic compromise (with consideration of pre-existing conditions).
- e. Abnormal respiratory rate not responding to initial treatment (with consideration of pre-existing conditions).
- f. Airway problems either before or after interventions.
- g. Signs/symptoms of hypoperfusion not improving.
- h. Decreased motor or sensory ability (with consideration of pre-existing conditions)

- i. Changes (deterioration) in presenting symptoms; stable patient who becomes unstable at any time.
- j. Consent problems and ALS Refusals.
- k. Uncertain triage decisions.
- l. Patients with a pulse in which transcutaneous pacemaker or electrical conversion therapy is used.

3. **COURTESY NOTIFICATION (CN):**

Required contact with receiving facility after ALS care according to treatment guidelines and reassessment. Vital signs are within normal limits, the patient's condition is stable or improved. No medical control input is required in addition to that covered under the treatment guidelines. This call is abbreviated and is designed to allow receiving facility to prepare for arrival.

**The following minimum information should be given during a "CN":**

- a. Identify self and agency
- b. Mechanism of injury
- c. Patient age, sex, chief complaint, vital signs, GCS, blood glucose and pertinent findings.
- d. Interventions, patient response/status
- e. ETA to hospital

4. **PATCH:**

Required on-line medical direction with Base Hospital (or back-up) (requires physician input). A patch includes the above information and a request for recommendations or general or specific treatment advice either from physician or his/her representative.

**EXCEPTIONS: (Critical Trauma, Medical Codes)**

In order to concentrate efforts on administering patient care and enhancing early communication to and preparedness of the receiving facilities of critical trauma patients and patients in cardiopulmonary arrest from medical causes, an abbreviated Courtesy Notification may be made with the receiving facility of these patients rather than a Patch under the following circumstances:

1. Appropriate treatment interventions are covered under trauma treatment guidelines and/or cardiopulmonary arrest treatment guidelines.
2. No question exists in the prehospital provider's judgment as to the Application/provision of care outlined in the specific Treatment Guidelines.
3. No additional medical direction is necessary in the prehospital provider's judgment for the provision of care and/or triage.

## **BASIC RADIO PROCEDURES**

All communications must include the following information:

1. EMSCOM Vehicle I.D.
2. Medic name & certification level
3. Status of call (A.L.S vs. B.L.S.)(Patch vs. Courtesy Notification)
4. Number of patients (If more than one patient)
5. Age & sex of patient(s)
6. Chief complaint(s)
7. History and objective finding(s)
8. Treatment rendered & response to treatment
9. State the orders you are requesting
10. E.T.A. and destination

## **COMMUNICATION GUIDELINES**

1. When using a radio, allow for a three-second delay after depressing the transmit key. This allows the electronics to fully engage.
2. Stop frequently and release transmit key to insure that the base hospital has received your transmission.
3. Ask for On-line Physicians to come on the line for any A.L.S. calls regarding patients you think might be unstable; or any time the scope of complexity of information requires direct contact with the physician.
4. Present information so that the listener gets an overview early (e.g. "... a 68 year old male, auto accident victim in acute respiratory distress..."). Report findings in the same order you evaluate a patient, i.e. primary assessment, vital signs, secondary assessment.
5. You need not list all relatively minor findings that do not affect immediate patient care decisions
6. Communicate with courtesy, brevity, and clarity.
7. Repeat all orders received back to the base hospital—medicine, dose, route, frequency.
8. Remember that many people are listening to your radio communications, do not use patient names and avoid use of unprofessional comments.
9. Follow Arizona Department of Public Safety (A.D.P.S.) EMSCOM Operations Manual.
10. Patches on B.L.S. patients should consume a minimum amount of time and only the most pertinent information.

## **COMMUNICATIONS SYSTEMS FAILURES**

If unable to contact the Base Station via Hospital Radio or dedicated phone lines, contact should be made with your alternate Base Hospital. Any situation where procedures are performed, which by these treatment guidelines require a medical control option, and such medical control option is not obtained because of failure to establish radio contact, will be reviewed individually as to their appropriateness. Clear cut indications for procedures must exist.

Base Hospitals shall develop plans for medical control in the event of local equipment failure. Such plans should include contingencies for radio failure, power outages, structural failures, etc.

## **INTERMEDIARY'S RESPONSIBILITY IN RADIO COMMUNICATION**

An intermediary is an emergency department nurse designated by the emergency physician to provide on-line medical supervision under verbal direction and control of the physician.

1. An intermediary will participate in daily communications and recording equipment troubleshooting procedure as outlined by A.D.P.S. R.C.C. Center policy.
2. An intermediary in contact with an A.L.S. unit will ask the emergency physician to come on-line at once if requested by the A.L.S. unit.
3. Communications with A.L.S. providers shall be completed in a timely, organized manner.
4. When a patient is to be transported to another receiving facility, immediately communicate all pertinent patient management information to the responsible physician or nurse at the receiving facility. If the receiving facility is also a Certified Base Hospital, direct communications with the A.L.S. unit rendering that care may be transferred to the receiving medical control authority at the discretion of the sending medical control authority, and with the knowledge and consent of the receiving medical control authority.
5. When relaying verbal directions/orders to field units, the intermediary shall identify by name the On-line Physician giving the orders transmitted.

## **BODY SUBSTANCE ISOLATION**

All patients should be considered potentially infectious. Standard precautions should be followed in accordance with Center for Disease Control (C.D.C.), Occupational Safety and Health Administration (O.S.H.A.), and base hospital guidelines.

## **TRANSPORTATION**

The patient should go to the medical facility which best meets his medical needs. If not the closest hospital, this decision requires a medical control option unless previously approved by the Administrative Medical Director. The patient's choice of hospital should be considered when such a request does not adversely affect or delay care or the operation of the transporting agencies.

If immediate hospital (medical/surgical) intervention is required, the quickest form of transport must be considered.

Scoop and Run involves rapid initiation of transport. It should not be undertaken until simple measures of airway control are performed on scene. The implementation of field procedures should not delay the transport of critical patients.

## **INTERFACILITY PATIENT TRANSPORTATION**

Interhospital patient transfers on an emergency basis are commonly initiated when definitive or therapeutic needs of a patient are beyond the capacity of one hospital. A pre patch needs to be made to the On-line Medical Direction Physician prior to leaving the sending facility with an ALS patient. Any change in patient status requires the personnel to contact their Base Hospital, not the receiving facility for further orders.

1. All patients should be stabilized as much as possible before transfer.
2. E.M.S. personnel must receive an adequate summary of the patient's condition, current treatment, possible complications, other pertinent information, and sending physician's determination of level of service needed during the transport.
3. E.M.S. ALS personnel continue to operate under control of the Base Hospital. Any orders given to such medics on interfacility transfers must be in accordance with their treatment guidelines and must be reviewed and approved by on-line medical control as the treatment guidelines specifies prior to transport.
4. Transfer papers, summary, lab work, X-rays, etc., should be given to the transporting E.M.S. personnel, not the family or friends.
5. The receiving hospital physician must be contacted by the transferring physician and agree to accept the patient prior to the transfer.
6. The level of emergency personnel must be appropriate to the treatment needed or anticipated during transfer.
7. Patients with intravenous infusion must be transported by the appropriate level of personnel. If a patient is receiving medication outside the scope of the transferring A.L.S provider, that patient must be accompanied by an R.N. or Physician as indicated by the patient's condition.

## **AT SCENE TRANSFER OF CARE/MULTIAGENCY DOCUMENTATION**

It is common for a variety of certified personnel with different skill levels to be providing care at the scene at one time. The fact that there is a higher skill level provider at the scene does not absolve each team member in patient care responsibilities.

Once on scene patient care is completed, and transportation of the patient is necessary, a few rules exist.

1. The A.L.S. provider with the highest skill level must accompany that patient to the receiving facility.
2. If care of the patient is transferred to another provider (that did not initiate the care), a report concerning patient scene, status, and care must be given to the provider when he or she accepts the patient.
3. Upon transfer of patient care, pertinent field information should be relayed without unnecessarily delaying transport.
4. When multiple teams render care, and do not arrive at the scene simultaneously, each team shall be responsible for reporting the care they rendered in written form. This is a minimum requirement and complex cases may require reporting of contemporaneous care

in multiple reports if the report writer was not able to keep adequate records during the call.

## **TRAPPED OR IMPALED PATIENT**

If you arrive at the scene to find a trapped or impaled patient who will take a significant time to extricate, or the impaled object cannot be easily cut, stabilize A.B.C.'s as much as possible and contact your Base Hospital. After explaining the situation, it may be appropriate for a physician from the hospital to come to the scene in case of the need for A.L.S. beyond your skill.

## **REFUSAL OF TREATMENT AND/OR TRANSPORT**

Every patient has the right to refuse treatment and/or transport. However, for a patient to be able to refuse treatment and/or transport the following criteria must be present:

- 1) Legal Competence (age 18 or emancipated minor)
- 2) Mental Competence (alert and oriented x 4)
- 3) Medical Competence (they must be able to clearly understand the medical consequences/ health risks for refusing treatment and transport)

VVMC does not support, condone, or allow EMS initiated refusal of transport. All refusals must be initiated by the patient or their guardian.

All patient refusals that involve ALS complaints and care require a patch to the base station. Medical control has the option to allow the refusal or to request the patient be restrained and brought to the ED for evaluation. This should only be done if it does not endanger the providers.

The patch for refusals should include the following:

- 1) Patient's chief complaint
- 2) 2 sets of vital signs (if able to obtain) \*\*\*the terminology "vital signs stable" is not acceptable, the patch must include the actual vital signs
- 3) Patient's physical exam
- 4) The patient's reason for refusal
- 5) Details on how the patient demonstrates legal, mental, and medical competence
- 6) The patient's plan for care or further evaluation

Documentation should include all of the above listed information required for the patch. It should also include any extra efforts done by providers (waiting on scene for parents/family to arrive, discussions with other persons on scene, obtaining phone numbers for call back, etc.)

BLS refusals should be documented identically to ALS refusals. These refusals do not require a patch, however providers are encouraged to patch if any unusual circumstances exist.

## **FIELD TRIAGE GUIDELINES**

Due to the rural and isolated nature of much of this region, coupled with the long distances between communities, the emergency patient is usually taken to the nearest Emergency Receiving Facility.

Exceptions may occur when:

1. A rational and oriented patient specifically requests transport to another facility, and the E.M.S. personnel deem it feasible to do so. This requires a medical control option. Specific agency policy may affect the decision.
2. The nature of the patient's illness or injury requires services not available at the nearest facility. The decision to bypass the nearest facility should be substantiated during direct communication with the responsible On-line Physician at the Base Hospital and in compliance with VVEMS Medical Direction Policy on Transport Destination.
3. Multiple victims have been identified by prehospital personnel and possible overloading of the nearest hospital's resources may prompt directing transport of a victim(s) directly to another facility.

Ordinarily, priority will be given to the most critical patients. However, when the number of patients exceeds the E.M.S. resources immediately available, then priority must be given to more salvageable patients.

## **MULTIPLE CASUALTY INCIDENTS (M.C.I.)**

If an agency has no formalized (written and implemented) M.C.I. Plan the following will briefly outline steps to be taken in the event of an M.C.I.

Definition of an M.C.I.:

1. Five (5) or more critically (Immediate) injured patients and/or
2. An incident that exceeds or potentially exceeds the E.M.S. resources available.

These are based upon common triage treatment guidelines and the use of a nationally recognized Incident Management or Command System (I.M.S. /I.C.S.). All agencies are expected to use the I.M.S. to allow agencies to work with a common system to mitigate incidents. This outline is not intended to replace well established local plans; rather, it offers a guideline for those areas in which no organized plan exists.

On arrival at an M.C.I. - in order of priority:

1. Perform scene size up, assure scene safety
2. Request additional resources:
  - a. from your agency;
  - b. Consider:
    - (1) Appropriate Law Enforcement Agencies
    - (2) Aircraft assistance
    - (3) Mutual aid
    - (4) Specialized needs (i.e. HazMat, School buses, etc.).
3. Establish Initial Command
4. Notify the Base Hospital that you have an M.C.I.
  - a. Number of patients

- b. Have Base Hospital notify regional hospital
- 5. When additional resources become available:
  - a. Assign per I.C.S. (i.e. Triage, Transportation, Staging, Safety, etc.).
  - b. START/Triage patients
    - 1. Immediate (to be transported first and treated immediately).
      - a. Respiration-over 30
      - b. Pulse-No Radial Pulse
      - c. Mental Status-Unable To Follow Simple Commands
    - 2. Delayed (transportation and treatment may be deferred).
      - a. Other patients unable to walk on their own
    - 3. Minor (to be transported or treated last)
      - a. Patients that can walk on their own.
    - 4. Dead/Dying
      - a. No Respirations after Head Tilt/OPA
  - c. Provide for scene security:
    - Safety officer/sector\* Law enforcement
  - d. Incident Command or Medical Group/Branch notifies receiving hospital of the number of patients and their categories.  
Additional contact should be made to the receiving hospital if there is a significant change in the number of patients they will be receiving.
- 6. Designate treatment areas for Immediate, Minor, and Delayed:
  - a. Mark areas with flags or tape with color designation
  - b. Move patients to proper treatment area.
  - c. Leave Dead/Dying victims where they are unless hindering other patient care
  - d. Treat patients in designated treatment area.
- 7. Transportation officer organizes transportation taking into consideration patient priority.
  - a. Transportation of patients to appropriate receiving facility(s)
  - b. Ensures adequate medical personnel remain on scene to treat remaining patients.
- 8. Ambulances will provide brief courtesy notifications to the receiving facility to include:
  - a. Triage priority of patients
  - b. Description of major injuries
  - c. Treatments provided
- 9. Consider Rescuer Assistance/Relief if incidents of long duration ("Rehab sector").
  - a. Arrange for food and water.
  - b. Rest area away from scene, if possible. (Consider house, store, etc.)
  - c. Rotate personnel through "Rehab Sector".
- 10. At conclusion of incident:
  - a. Restock units
  - b. Consider post incident debriefing for all Rescuers and Police.
    - (1) Within 12 hours post-incident.
    - (2) Follow-up within 72 hours.
    - (3) Offer individual counseling if needed/available.

Note: The above does not offer a detailed, in-depth study of M.C.I. response or the I.C.S. system. Further education in these areas should be pursued as space here will not allow total coverage of these areas. Practical drills and daily use of the I.C.S. on all multi-casualty incidents will increase proficiency in these areas.

## **PNEUMATIC ANTI-SHOCK GARMENT (P.A.S.G.)**

The PASG has a limited role in modern EMS care. ADHS no longer requires PASG to be present on EMS vehicles and we only support its use in stabilizing pelvic fractures when other services are impractical or unavailable.

## **TREATMENT GUIDELINES**

### **GENERAL ASSESSMENT AND TREATMENT APPROACH**

Although there are many things that may be medically affecting your patient, there are a limited number of supporting treatments you have to offer. Do not let the gathering of information distract you from the management of life-threatening problems.

Remember, however that you may be able to gather information from bystanders at the scene, from the environment, and perhaps even from the patient that may not be available to the physician later on. Your partner can often be engaged in collecting this kind of information during the secondary examination.

### **HISTORY**

1. Chief complaint (questioning to include, when appropriate):
  - a. Onset
  - b. Provocation
  - c. Quality
  - d. Radiation
  - e. Severity
  - f. Time
2. Associated complaints: question as for Chief complaint
3. Relevant past medical history
4. Allergies
5. Medications and drugs: chronic
6. Survey of surroundings for evidence of drug abuse, mental functioning, and family problems
7. Last meal, last menstrual period (if applicable)

### **INITIAL ASSESSMENT**

Primary interventions should always be made as soon as a need for them is assessed.

#### **AIRWAY:**

Assess patency, stridor, foreign body (F.B.), ability to maintain airway.

### **TREATMENT**

1. If compromised or absent airway, or patient unresponsive:
  - a) Position the airway
  - b) Insert OPA/NPA
  - c) Suction PRN
  - d) Remove dentures

- e) Always consider C-spine injury
2. Consider endotracheal intubation or Combitube
3. Consider needle or surgical cricothyrotomy

### **BREATHING:**

Assess: Rate, apparent tidal volume, effort, ability to speak, symmetrical movement, breath sounds, accessory muscle use, oximetry.

Realize that oxygenation and ventilation are separate but interdependent issues. Oxygenation may be assessed as adequate with a pulse oximeter, but the only way to assess ventilation as adequate is by ETCO<sub>2</sub> monitoring and/or clinical means, i.e. rate, tidal volume, air movement.

### **TREATMENT**

1. Position of comfort when appropriate
2. Oxygen as appropriate
3. Assist with Bag-Valve mask
4. CPAP may be used when indicated by protocols.

### **CIRCULATION:**

Assess pulse presence, location, quality, and capillary refill; assess blood loss from hemorrhage, skin color and temperature, and level of consciousness.

### **TREATMENT**

1. Control active external bleeding with direct pressure, splint major fractures
2. IV NS; consider volume support (enroute)
3. Monitor Rhythm
4. Drug therapy as indicated

### **VITAL SIGNS**

1. Obtain first quantitative set of vitals within five minutes if practical (pulse, blood pressure, respiratory rate, pulse oximetry, temperature)
2. Repeat according to patient's condition. At least one more set prior to transport or enroute.

### **NEUROLOGICAL ASSESSMENT**

Management of patients with head injury or neurological illness depends on careful assessment of neurological function. Changes in neurologic status are particularly important. The first observation of neurological status in the field provides the basis for monitoring sequential changes. It is, therefore, important that the first responder accurately observe and record neurological assessment, using parameters which will be followed throughout the patient's hospital course.

- The Glasgow Coma Scale is one method of monitoring patients with head injury. Errors and confusion are minimized when precise responses to specific stimuli are recorded. Always record specific responses in addition to the total score of the Glasgow Coma Scale. See Appendix E for Glasgow Coma Scale
- Another method to objectively describe LOC in the non-head injured patient is **AVPU**
  - A:** Awake & Alert
  - V:** Responsive to Verbal Stimulus
  - P:** Responsive to Painful Stimulus

**U: Unresponsive**

- Eyes:
  1. Direction of gaze
  2. Size and reactivity of pupils
  3. Visual Field Loss
- Motor Function and Coordination
  1. Observe whether all four extremities move equally well
  2. Facial Droop
- Speech and Language
  1. Real words, but slurred enunciation
  2. Unable to use correct words and/or unable to comprehend simple question and commands
- Sensation (if patient awake):
  1. Observe for absent, abnormal or normal sensation at different levels if cord injury is suspected

**SPECIAL NOTES:**

- A. Sensory and motor exam **must** be documented before and after moving patient with suspected spinal injury.
- B. Note what stimulus is being used when recording responses.

**GENERAL: FOCUSED HISTORY/PHYSICAL EXAM OR RAPID ASSESSMENT**

**DETAILED PHYSICAL EXAM Definitions:**

Focused History/Physical Exam: The part of the assessment process in which the patient's major complaints or any problems that are immediately evident are further and more specifically evaluated.

Detailed Physical Exam: The part of the assessment process in which a detailed area-by area exam is performed on patients whose problems cannot be readily identified or when more specific information about problems identified in the focused history and physical exam is necessary.

The four components of physical examination are: inspection, auscultation, palpation, and occasionally, percussion.

See Appendix H for Detailed Head to Toe Assessment

**CENTRAL VENOUS ACCESS**

Administrative Medical Control has not authorized the initiation of central venous lines by Paramedics. Existing central venous access devices such as, porta-caths and PICCs may be accessed by Paramedics only as trained in the Advanced IV access training.

## **CPAP**

CPAP, an optional respiratory support treatment that has shown to rapidly improve vital signs, gas exchange, and work of breathing shortness of breath, decrease the sense of dyspnea, and decrease the need for endotracheal intubation in patients who suffer from asthma, COPD, pulmonary edema, CHF, and pneumonia. This is approved for use by VVEMS agencies after proper training as delineated in the CPAP Use Guideline. See Appendix G


## **Intraosseus IV Access/IO**

VVEMS agencies may use this technique as part of their vascular access after completing the required training, in accordance with the IO Use Guideline. See Appendix G

## **Rapid Sequence Intubation/RSI/Medication Assisted Intubation**


This is an approved skill for EMS use in the state of AZ. Only agencies currently using RSI are approved use at VVMC. VVEMS medical direction supports the use of RSI as an optional advanced airway

# DEAD ON ARRIVAL



1) In situations where hypothermia may be a consideration, hypothermia guidelines should be followed. Seek Medical Control input.


# DO NOT ATTEMPT RESUSCITATION ORDERS



- (1) It is not the intent of advanced directives to deny treatment of other medical conditions not related to the terminal illness, pain medication, or other supportive care.
- (2) If patients relatives are present and are indicating they want resuscitation attempted, in the presence of advance directives, begin resuscitation and patch for Medical Control input.
- (3) If patient is in a healthcare facility or is being transported interfacility with a physician's DNAR in place it is not necessary to begin CPR.

# WITHHOLDING/TERMINATION OF RESUSCITATION

I-99 Guideline see Appendix K




(1) For indications where no treatment is required the provider may withhold resuscitative efforts and patch for medical direction.  
(2) Consideration should be given to potentially reversible conditions such as, overdose/poisoning, hypothermia, cold water drowning, etc.

# ADULT BRADYCARDIA, UNSTABLE (1)

I-99 Guideline see Appendix K

HEART RATE < 60 MINUTE WITH ACCOMPANYING SIGNS/SYMPTOMS OF HEMODYNAMIC COMPROMISE, I.E., CHEST PAIN, HYPOTENSION, IF HISTORY/EVIDENCE OF TRAUMA, PROCEED TO TRAUMA TREATMENT GUIDELINE




- (1) Signs/symptoms of an unstable patient may include chest pain, SOB, decreased LOC, hypotension, shock, pulmonary edema, congestive heart failure, and acute myocardial infarction.
- (2) This should not delay definitive treatment.
- (3) Repeat vital signs and lung auscultation before and after fluid administration
- (4) Contact Medical Control to consider administration of Dopamine 5-20 mcg/kg/min and/or Epinephrine 2-10 mcg/min

## CHEST PAIN SUGGESTIVE OF CARDIAC ORIGIN


Chest Pain suggestive of possible myocardial ischemia (1)

I-99 Guideline see Appendix K



- 1) Indications of chest pain suggestive of possible myocardial ischemia include: Description of crushing, squeezing, pressure, burning, tightness, diaphoresis, nausea/vomiting, apprehension, radiation, age>30, **associated cardiac risk factors**.
- 2) If twelve lead capability, should be done in pt's initial assessment.
- 3) Repeat vital signs and lung auscultation before and after administration of NTG. Consider prior NTG use. If pain reoccurs and is not refractory to NTG, repeat NTG 0.4mg SL every 5 minutes as needed for pain relief, maintaining B/P > 100.
- 4) Nitroglycerin is contraindicated in patients that have taken Viagra (sildenafil), Cialis, Levitra, or similar medications in the previous 72 hours
- 5) Contraindication to Aspirin if has an allergy to ASA. Can administer ASA without and IV in place.
- 6) Communication with hospital should be completed as soon as possible so that Cath Lab team can be notified for ST Elevation MI.


## ADULT CARDIOPULMONARY ARREST – CCR ALTERNATIVE (1)(2)



- 1) Age greater than 8 years old.  
 2) Should not be used on patients; involved in traumatic event, overdose or where evidence of primary respiratory arrest is present.  
 3) Do not attempt intubation until after 4<sup>th</sup> set of 200 chest compressions. Ventilate with BVM if necessary.  
 4) Pulse checks should be done only if ECG indicates a potentially perfusing rhythm. Do not interrupt chest compressions. Be very brief.

## ADULT PULSELESS ARREST


I-99 Guideline see Appendix K



- 1) Assess rhythm – quick look, only check pulses if there is an organized rhythm present.
- 2) Evaluate airway, intubate if necessary, limit interruption of CPR as much as possible.
- 3) Once patient is successfully intubated perform continuous asynchronous compression (rate 100/min) with ventilations (rate 8-10/min)
- 4) Pulse checks should be done only if EKG indicates a potentially perfusing rhythm, do not interrupt chest compressions, and be very brief.
- 5) Medications should be administered during CPR as soon as possible after rhythm checks.
- 6) Consider possible causes: Hypovolemia, (volume infusion), hypoxia (ventilation/re-evaluation), acidosis (ventilation/re-evaluation), tension pneumothorax (needle decompression), hypothermia, hypoglycemia, drug overdose, cardiac tamponade (volume infusion), massive AMI, hyperkalemia (consider NaHCO<sub>3</sub>, D50W, Calcium Chloride) massive pulmonary embolism.
- 7) If patient remains asystolic or other agonal rhythm after successful intubation, initial medications, no reversible causes are identified, and transport has not been initiated, consider termination of resuscitative efforts by order of a physician. Consider interval since arrest.
- 8) For successful conversions with HR>60 and no 2<sup>nd</sup> or 3<sup>rd</sup> degree heart blocks. Assess vital signs, administer Lidocaine 1-1.5 mg/kg and start infusion at 2-4 mg/min. or Amiodarone 150 mg IV over 10 minutes then begin drip at 1mg/min for first 6 hours. If patient received bolus doses prior to conversion administer maintenance infusion only.

# CARDIAC ARREST POST RESUSCITATION INDUCED HYPOTHERMIA

I-99 Guideline see Appendix K



1) Criteria for Induced Hypothermia:

- ROSC after cardiac arrest not related to trauma or hemorrhage
- Age greater than 18 years old
- Patient is intubated and remains comatose (no purposeful response to pain)
- GCS of less than 8 after ROSC
- ROSC within 60 minutes of arrest
- Initial temperature >34C- no environmental hypothermia related arrhythmia
- Female patients not pregnant
- No uncontrolled hemorrhage
- No persistent unstable arrhythmia
- No DNR paperwork identified during resuscitation

2) If unable to intubate DO NOT initiate induced hypothermia

3) Patients develop metabolic alkalosis with cooling. Do not hyperventilate.

4) When exposing pts for cooling purposes undergarments may remain in place. Be mindful of your environment and be preserve patients modesty.


5) Do not delay transport for purposes of cooling.

6) Patch early to advise of induction of hypothermia to allow ED to have equipment ready.

7) **AT ANY TIME- Loss of Spontaneous Circulation- discontinue cooling and go to appropriate protocol**


# ADULT TACHYCARDIA WITH PULSES

I-99 Guideline see Appendix K




- 1) If at any time patient becomes unstable, proceed to "Unstable" side
- 2) Carotid sinus massage should not be performed without Medical Control Contact ; other methods of vagal stimulation should be attempted. Carotid sinus massage is contraindicated if patient >50 years of age or has history of hypertension. If ordered by Medical Control, verify absence of carotid bruits.
- 3) Contact Medical Control to administer Diltiazem (Cardizem) 0.25 mg/kg, if no response may repeat in 15 minutes at 0.35mg/kg. Amiodarone 150 mg administered over 10 minutes, if no response may repeat every 10 minutes, maintenance infusion after conversion is 1 mg/min. Consider cardioversion
- 4) For successful conversions of ventricular arrhythmias with HR > 60 and no 2<sup>nd</sup> or 3<sup>rd</sup> degree heart blocks: Assess vital signs, administer Amiodarone 150 mg IV over 10 minutes then begin drip at 1 mg/min. for first 6 hours or Lidocaine (Xylocaine) 1mg/kg and start infusion at 2-4 mg/min, reduce maintenance infusion of Lidocaine by half in patients with renal or hepatic disease or > 70 years of age. If patient received bolus doses prior to conversion administer maintenance infusion only.
- 5) If delays in synchronization occur or rhythm is polymorphic VT go immediately to unsynchronized defibrillation at 120-200 biphasic with manufactures recommendations or monophasic 360J. For polymorphic VT the provider should be prepared to move immediately to the Pulseless Arrest algorithm if pulseless arrest develops.

# TRAUMA TRIAGE DESIGNATION




## SUBMERSION INCIDENT




- 1) BVM with reservoir with 100% O<sub>2</sub> may be adequate to provide ventilation and oxygenation. If ventilation appears clinically inadequate or transport will be greater than 5 minutes, consider intubation.
- 2) 100% oxygen should be used in all patients.
- 3) Rapid transport is of the utmost importance. Advanced Life Support procedures should be attempted at the scene, but if unsuccessful within a short period of time, the patient should be transported to nearest appropriate facility without further delay.
- 4) Gastric decompression allows adequate pulmonary tidal volumes. Insert 16-18 FR. NG/OG catheter.
- 5) Establishment of an IV should not delay patient transport.
- 6) Hypoxia (ventilation/re-evaluation), acidosis (ventilation/re-evaluation, consider orders for sodium bicarbonate), tension pneumothorax (needle decompression), hypothermia (see Hypothermia Treatment Guideline), trauma-hypovolemia (volume infusion), hypoglycemia (check blood sugar)

## TRAUMA - BURNS




1) If patient or clothing still burning cool hot areas immediately. Flush chemical burns for 20 minutes.

## TRAUMA – MUSCULOSKELETAL INJURY



1) Consider Air Transport for isolated replantable extremities


## TRAUMA – HEAD INJURY WITH ALOC (1)



- 1) GCS 13 or less, consider Air Transport to Neurological Center. Discuss with patch MD
- 2) Controlled hyperventilation with 100% O<sub>2</sub> at 20 breaths per minute should only be used in patients with signs of impending central herniation; unconscious, unresponsive patient with extensor posturing or no motor response; asymmetric or dilated and unreactive pupils; GCS decreases 2 or more points from patient's prior best score when patient had initial GCS of 9 or less, after correction of hypoxemia, hypotension, and hypoglycemia. *Normoventilation is 10 bpm in the adult.*
- 3) The goal for time on scene is to not exceed ten (10) minutes for patient assessment, management and packaging unless extrication is required or unforeseen circumstances develop.
- 4) On-line Medical Control should be involved in difficult or questionable triage decisions.
- 5) Consider IO if no IV access and patient is in extremis.

## TRAUMA – MULTI – SYSTEM


Applies to patients presenting with S/S of Critical (Immediate) injury or patients in which the mechanism of injury is suspect for occult critical Injury.



- 1) The goal for time on scene is not to exceed ten (10) minutes for patient assessment, management and packaging unless extrication is required or unforeseen circumstances develop. Patients with penetrating injuries to the thorax or head with unstable vital signs should be transported immediately.
- 2) On-line Medical Control should be involved in difficult or questionable Triage decisions.
- 3) Consider IO if no IV access and patient is in extremis.
- 4) PASG/MAST is contraindicated in penetrating chest trauma and is relatively contraindicated in isolated blunt chest trauma.
- 5) Careful consideration should be given to the amount of fluids infused in the field.

# TRAUMA – SPINAL INJURY


I-99 Guideline see Appendix K



1) If patient remains hypotensive and appears to have isolated head / spinal injuries contact Medical Control to administer Dopamine 5-20 mcg/kg/min.

# CEREBRAL VASCULAR ACCIDENT – STROKE

I-99 Guideline see Appendix K




1) Establishing time signs and symptoms began is CRITICAL. If patient awoke from sleep with S/S it is also important to determine how long patient was asleep. Patients with ischemic strokes < 3 hours old may be candidates for TPA therapy with some candidates eligible for up to 4.5 hours.

2) Do not treat patients for hypertension unless directed to do so by medical control.

## ALLERGIC REACTION

Applies to patient presenting with systemic allergic reaction e.g. diffuse urticaria, angioedema (edema of deep dermis layers), abdominal cramping, nausea or vomiting without anaphylaxis




- 1) The use of Epinephrine in patients 45 years or greater with known coronary artery disease requires Medical Control input.  
2) Consider acuity of onset of symptoms and history of prior anaphylactic reaction.  
3) If IV cannot be established, administer Benadryl (Diphenhydramine) 50 mg IM.

## ANAPHYLAXIS

Applies to patient presenting with allergic reaction and with signs and symptoms of airway, respiratory, or circulatory compromise (laryngeal edema, bronchospasm, or hypotension.)


**I-99 Guideline see Appendix K**



- 1) If signs and symptoms of severe hypoperfusion and an IV can be rapidly established, consider going directly to IV Epinephrine as per protocol.
- 2) Establishment of an IV should not delay the administration of IM Epinephrine to a patient in extremis.
- 3) The use of Epinephrine in patients age > 45 years or with known coronary artery disease requires Medical Control input.
- 4) If prolonged transport consider repeat use of Epinephrine every 15 minutes. Medical Control input should be obtained, if possible.
- 5) At any time an IV cannot be established, give Benadryl ( Diphenhydramine) 50 mg IM as soon as possible after Epinephrine IM.
- 6) Consider IO if no IV access and patient is in extremis.
- 7) Consider the use of SVN therapy via in line BVM system in patients who are tiring or are appearing to have decreased tidal volumes.
- 8) If patient continues to be hypotensive contact Medical Control to administer Epinephrine 1:1,000 2-10 mcg/min IV/IO Infusion, titrate to effect or Dopamine drip 5-20 mcg/kg/min.


# ENVENOMATION – ARACHNIDS

I-99 Guideline see Appendix K



- (1) Attempts to kill or capture insect or bring to ED are not recommended.
- (2) Contact Medical Control to administer Versed ( Midazolam) / Valium (Diazepam) for severe pain / muscle spasm.
- (3) Careful observation of respiratory status.

## ENVENOMATION – SNAKE BITES




1) Attempts to kill or capture the snake or bring dead animal to ED are NOT recommended.

2) Many exotic snakes are neurotoxic so respiratory status must be monitored carefully.

3) Contact medical control for Epinephrine if patient is hypotensive. If presenting with allergic reaction follow Allergic Reaction protocol.

## AIRWAY COMPROMISE


I-99 Guideline see Appendix K



- 1) Medical Control contact is not mandatory, however, the medic is encouraged to discuss with Medical Control if he/she is anticipating a Cricothyroidotomy and the clinical situation is such that there is time for Medical Control contact.
- 2) Consider NG/OG tube for gastric decompression in situations of prolonged ventilation.
- 3) Consider blind airway insertion device if difficult airway and Endotracheal and Nasotracheal intubation unsuccessful after two attempts.

# AIRWAY OBSTRUCTED


I-99 Guideline see Appendix K



1) Chest thrusts if patient is obese or pregnant.  
2) Verify proper tube placement by bulb tube check / air aspiration or ETCO2 detector / monitor, chest wall rise, good breath sounds, absence of gastric sounds, and clinical improvement in patient.

## RESPIRATORY INSUFFICIENCY – BRONCHOSPASM


Applies to patients with S/S of acute respiratory distress, secondary to asthma, COPD, and inhalation injury  
**I-99 Guideline see Appendix K**



- 1) Administer O2 at high flow rates to all patients in severe respiratory distress. This is especially true if pulse oximetry is not available.
- 2) Consider the use of SVN therapy via in line BVM system in patients who are tiring or are appearing to have decreased tidal volumes.
- 3) The use of epinephrine in patients 45 years or greater or with known coronary artery disease requires **Medical Control** input.
- 4) Do not delay definitive therapy to establish IV.
- 5) Obtain an IO if no IV access and patient is in extremis.
- 6) Atrovent (Ipratropium Bromide) is contraindicated in patients with soy or nut allergy.
- 7) Epinephrine IM is indicated for use in bronchospasm i.e. bronchiolitis and asthma
- 8) Magnesium is for bronchospasm / asthma only NOT COPD


# RESPIRATORY INSUFFICIENCY – PULMONARY EDEMA

I-99 Guideline see Appendix K




- 1) Patients who appear to be tiring or have decreased tidal volume may require respiratory assist.
- 2) High flow O<sub>2</sub> should be used in any patient who appears distressed
- 3) Repeat vital signs and lung auscultation before and after administration of NTG.
- 4) Contact Medical Control to administer Dopamine drip 5-20mcg/kg/min.
- 5) Unless directed by online medical control Nitroglycerin is contraindicated in patients that have taken Viagra (sildenafil), Cialis, Levitra, or similar medication within 72 hours

**OBSTETRICS  
COMPLICATIONS OF DELIVERY  
ABNORMAL PRESENTATIONS**




**OBSTETRICS**  
**COMPLICATIONS OF DELIVERY**  
**POST PARTUM HEMORRHAGE (1)**




1) Post partum hemorrhage is defined as blood loss in excess of 500mL and during the first 24 hours after delivery.

## OBSTETRICS COMPLICATIONS OF PREGNANCY




1) Signs of PIH/ pre-eclampsia / eclampsia may include: Diastolic BP . 80 mmHg with cerebral or visual disturbances, epigastric or RUQ pain with nausea and vomiting, ALOC, hyper-reflexia, peripheral edema, pulmonary edema, seizures.

## OBSTETRICS COMPLICATIONS OF PREGNANCY CONT.



## OBSTETRICS DELIVERY




1) Prepare for immediate delivery if (a) contractions are less than 2 minutes apart and/or (b) perineal bulge obvious and scalp becomes visible (crowning)

## ALTERED LEVEL OF CONSCIOUSNESS

GCS of 14 or <, psychotic or combative behavior, the post seizure patient, the near/post syncope patient, or any patient with history of ALOC as a part of current event.(1)

**I-99 Guideline see Appendix K**




- 1) Utilize information obtained from family, bystanders, friends, or other health care workers.
- 2) If hypoglycemia or opiate OD suspected, BLS airway management maybe sufficient until response to Dextrose and/ or Naloxone is determined.
- 3) If no history of alcoholism is suspected and malnutrition or cachexia is not present, Thiamine may be withheld.
- 4) If no change in LOC, repeat glucose. Realize the onset of action of Glucagon is 5-15 minutes.
- 5) Consider 12 lead

## ALTERED LEVEL OF CONSCIOUSNESS

With suspected alcohol intoxication (1)


I-99 Guideline see Appendix K



- 1) Utilize information obtained from family, bystanders, friends, or other health care workers.
- 2) If hypoglycemia or opiate OD suspected, BLS airway management maybe sufficient until response to Dextrose and/ or Narcan ( Naloxone) is determined.
- 3) If no change in LOC, repeat glucose. Realize the onset of action of Glucagon is 5-15 minutes.
- 4) Consider 12 lead
- 5) Do not intubate sleeping, stable intoxicated patient if oxygenating and ventilating.

# ADULT VIOLENT/AGITATED PATIENT (1)


I-99 Guideline see Appendix K



- (1) Agitated patient (significant psychomotor agitation) due to possible drug ingestion and/or violent patients who after initial treatment remain a danger to self or others.
- (2) Patients may need to be restrained for patient, provider, or bystander safety in order to complete other appropriate treatment guidelines.
- (3) If patient is in police custody and handcuffs have been applied it is preferable that a police officer also accompany the patient. EMS providers must, at a minimum, have the handcuff key in their possession during transport.
- (4) Patients shall be positioned in a manner that does not compromise airway or breathing. No patient will be restrained prone or "hog-tied." No patient will be placed between backboards or gurneys.

# SEIZURE

Prolonged, Repetitive, or Status Epilepticus  
**I-99 Guideline see Appendix K**




1) If IEMT is highest level of care may give Valium (Diazepam) 0.1 mg/kg IV every 5 minutes prn maximum of three doses.  
 2) The only approved intranasal method of administering medications is with the Mucosal Atomizing Device (MAD). Administer no more than 1ml per nostril. Use higher concentration of Versed 5mg/ml.


# ABDOMINAL PAIN, NON – TRAUMATIC

Testicular torsion, Pelvic pain, AAA

I-99 Guideline see Appendix K




# NAUSEA AND VOMITING




## ENVIRONMENTAL – HEAT RELATED

I-99 Guideline see Appendix K



- 1) Do not cool below 102 degrees F.  
 2) Do not over cool and cause shivering and reoccurring heat buildup. If patient is shivering contact Medical Control to administer Midazolam or Diazepam.  
 3) If patient is agitated contact Medical Control to administer Midazolam or Diazepam.

## ENVIRONMENTAL - HYPOTHERMIA




- 1) If there is an organized rhythm do not begin CPR unless directed by Medical Control.  
 2) Utilize only 1 shock.  
 3) Contact Medical Control for ACLS medication administration regimen. Consider withholding medications if core temperature is  $\leq 86$  degrees F and an extended time between doses if temperature is  $> 86$  degrees F.

## HYPOTENSION, NOT TRAUMATIC (1)


Applies ONLY when other specific ALS protocols do not apply.  
Hypotension is defined as BP < 90 systolic and associated signs / symptoms of hypoperfusion.  
If history / evidence of Trauma, proceed to Trauma Treatment Guideline.

### I-99 Guideline see Appendix K



- 1) PMH and patient's medications may be key to index of suspicion for cause of hypotension, e.g. history of ulcers, aneurysm, previous cardiac disease, alcoholism, etc. Consider possible causes of hypotension and treat cause.
- 2) Consider establishing 2 large bore IV's dependent upon patient's presentation
- 3) Bolus fluid in less than 10 minutes.
- 4) Repeat vital signs and lung auscultation before and after fluid administration.
- 5) If pulsatile abdominal mass present or suspected AAA/TAA, PATCH.
- 6) If patient continues to be hypotensive contact Medical Control to administer Dopamine drip 5-20 mcg/kg/min.


## POISONING / OVERDOSE (1)



- 1) Patients who are suspected or known to have ingested substances with a suicidal intent may refuse transport.
- 2) Bring bottles / containers if possible. INSPECT SCENE.
- 3) Consider Medical Control input for Sodium Bicarbonate 1-2 mEq/kg for TCA overdose, Calcium chloride 0.5 -1 Gm. for calcium channel blocker overdose, Atropine 2 mg every 2-4 min. for organophosphate exposure.
- 4) Do not intubate a stable airway. Give O2 and Ventilate patient.

## PEDIATRIC BRADYCARDIA, UNSTABLE


I-99 Guideline see Appendix K



- 1) If airway is managed with BVM for greater than 2 minutes, insert 10-16 Fr. OG/NG tube. Gastric decompression allows adequate pulmonary tidal volumes.
- 2) Special considerations may apply in the presence of severe hypothermia.
- 3) Consider IO use if IV access unavailable.
- 4) Dilute 1:1000 Epinephrine with 3-5 ml of NS flush.
- 5) Limited pediatric data; 15 kg or less pediatric electrodes recommended. For greater than 15 kg use adult electrodes.
- 6) Consider Medical Control input to administer Epinephrine IV continuous infusion at a rate of 0.1 to 1 mcg/kg/min.
- 7) rapid transport is essential in these situations. The above procedures should be performed as the patient is being moved towards the hospital.

# PEDIATRIC PULSELESS ARREST


I-99 Guideline see Appendix K



- 1) Assess rhythm – quick look, only check pulses if there is an organized rhythm present.
- 2) Evaluate airway, intubate if necessary, limit interruption of CPR as much as possible.
- 3) Once patient is successfully intubated perform continuous asynchronous compression (rate 100/min) with ventilations (rate 8-10/min)
- 4) Pulse checks should be done only if EKG indicates a potentially perfusing rhythm, do not interrupt chest compressions, and be very brief.
- 5) Medications should be administered during CPR as soon as possible after rhythm checks.
- 6) Consider possible causes: Hypovolemia,(volume infusion), hypoxia (ventilation/re-evaluation),acidosis (ventilation/re-evaluation), tension pneumothorax (needle decompression), hypothermia, hypoglycemia, drug overdose, cardiac tamponade (volume infusion), massive AMI, hyperkalemia (consider NaHCO<sub>3</sub>, D50W, Calcium Chloride) massive pulmonary embolism.
- 7) If airway managed with BVM > 2 min. insert 10-16 Fr. OG/NG tube after patient has been intubated.
- 8) If patient remains asystolic or other agonal rhythm after successful intubation, initial medications, no reversible causes are identified, and transport has not been initiated, consider termination of resuscitative efforts by order of a physician. Consider interval since arrest.

# PEDIATRIC TACHYCARDIA WITH PULSES


I-99 Guideline see Appendix K



- 1) Probable SVT in pediatrics: History incompatible with presentation, P waves absent or abnormal, HR not variable with activity, abrupt rate changes. Infant rate usually > 220 bpm and children usually > 180 bpm.
- 2) Patients often fit in between borderline and critically unstable situations. In these circumstances, a trial of adenosine may be considered but the medic must be prepared for immediate cardioversion.
- 3) the medic should consult Medical Control and consider reducing the Adenosine dosage in patients who are on Dipyridamole (Persantine) and Carbamazepine ( Tegretol)
4. Consider 12 lead EKG
- 5) Or biphasic equivalent
- 6) If probable VT contact Medical Control to administer Amiodarone 5mg/kg, max single dose 150 mg over 20 minutes may repeat two more times to a total of 15 mg/kg/day or Lidocaine 1mg/kg every 5-10 minutes to a total of 3 mg/kg.

# PEDIATRIC – NEONATAL RESUSCITATION

I-99 Guideline see Appendix K




- 1) If patient is not vigorous and meconium staining is present deep suction mouth and posterior pharynx then nose. Tracheal suctioning may be necessary before stimulating neonate and proceeding with other resuscitative steps. Vigorous- strong respiratory effort, good muscle tone, heart rate > 100 bpm. Depressed- weak or absent respiratory effort, poor muscle tone/limp, heart rate < 100 bpm.
- 2) Tracheal intubation may be considered at several steps. Tracheal tube should be used for tracheal suctioning.
- 3) Utilize IO or if peripheral IV sites inaccessible.

## PEDIATRIC – SUBMERSION INCIDENT – CATEGORY 1

Applies to a patient with no spontaneous respirations or pulses on arrival of unit; also includes patient with pulses and respirations and with significant alteration of LOC.


### I-99 Guideline see Appendix K



- 1) BVM with reservoir with 100% O<sub>2</sub> may be adequate to provide ventilation and oxygenation. If ventilation appears clinically inadequate or transport will be greater than 5 minutes, consider intubation.
- 2) 100% oxygen should be used in all patients.
- 3) Rapid transport is of the utmost importance. Advanced Life Support procedures should be attempted at the scene, but if unsuccessful within a short period of time, the patient should be transported to nearest appropriate facility without further delay.
- 4) Gastric decompression allows adequate pulmonary tidal volumes. Insert 10-16 Fr. NG/OG catheter.
- 5) Establishment of an IV should not delay patient transport.
- 6) Hypoxia (ventilation/re-evaluation), acidosis (ventilation/re-evaluation, consider orders for sodium bicarbonate), tension pneumothorax (needle decompression), hypothermia (see Hypothermia Treatment Guideline), trauma-hypovolemia (volume infusion), hypoglycemia (check blood sugar)


## PEDIATRIC – SUBMERSION INCIDENT – CATEGORY 2

Applies to a patient presenting alert and oriented (may have a history of altered level of consciousness prior to arrival of the rescue unit ) with spontaneous respirations and heart rate. (1)



1) These children require further medical evaluation. Child should be transported via ALS (if available) ambulance to the closest emergency care facility.


## PEDIATRIC TRAUMA - BURNS



1) If patient or clothing still burning cool hot areas immediately. Flush chemical burns for 20 minutes.


# PEDIATRIC TRAUMA – MUSCULOSKELETAL INJURY

I-99 Guideline see Appendix K



1) Patients under 15 years of age.  
2) Contact Medical Control to administer Midazolam 0.05mg/kg or Diazepam 0.1mg/kg for large muscle spasm.


## PEDIATRIC TRAUMA – HEAD INJURY WITH ALOC (1)



- 1) GCS less than or equal to 13 , consider Air Transport to Pediatric Neurological Center.
- 2) Minimize attempts or risks if patient is stable. Controlled hyperventilation with 100% O<sub>2</sub> at 20 breaths per minute should only be used in patients with signs of impending central herniation; unconscious, unresponsive patient with extensor posturing or no motor response; asymmetric or dilated and unreactive pupils; GCS decreases 2 or more points from patient's prior best score when patient had initial GCS of 9 or less, after correction of hypoxemia, hypotension, and hypoglycemia. *Normoventilation is 10 bpm in the adult.*
- 3) The goal for time on scene is to not exceed ten (10) minutes for patient assessment, management and packaging unless extrication is required or unforeseen circumstances develop.
- 4) On-line Medical Control should be involved in difficult or questionable triage decisions.
- 5) Consider IO if no IV access and patient is in extremis.

## PEDIATRIC TRAUMA – MULTI – SYSTEM


Applies to patients presenting with S/S of Critical (Immediate) injury or patients in which the mechanism of injury is suspect for occult Critical injury.



1) OG/NG tube if child ventilated with BVM for > 2 minutes or obvious gastric distention.  
 2) The goal for time on scene is not to exceed ten (10) minutes for patient assessment, management and packaging unless extrication is required or unforeseen circumstances develop.  
 3) Careful consideration should be given to the amount of fluids infused in the field.  
 4) On-line Medical Control should be involved in difficult or questionable Triage decisions.


# PEDIATRIC TRAUMA – SPINAL INJURY

I-99 Guideline see Appendix K



1) If patient remains hypotensive and appears to have isolated head / spinal injuries contact Medical Control to administer Dopamine 5-20 mcg/kg/min.

## PEDIATRIC ALLERGIC REACTION




1) Consider acuity of onset of symptoms and history of prior anaphylactic reaction.

2) If IV cannot be established administer Benadryl (Diphenhydramine) 1 mg/kg up to 25 mg IM as soon as possible after Epinephrine IM.

## PEDIATRIC – ANAPHYLAXIS


Applies to patient presenting with allergic reaction and with signs and symptoms of airway, respiratory, or circulatory compromise (laryngeal edema, bronchospasm, or hypotension).



- 1) If prolonged transport, consider repeat Epinephrine every 10-15 minutes. Medical Control input should be obtained, if possible.  
 2) Establishment of an IV should not delay the administration of Epinephrine IM to a patient in extremis.  
 3) At any time an IV cannot be established, give Benadryl ( Diphenhydramine) 1 mg/kg up to 25 mg IM as soon as possible after Epinephrine IM.


## PEDIATRIC ENVENOMATION – ARACHNIDS

I-99 Guideline see Appendix K




- 1) Attempts to kill or capture insect or bring to ED are not recommended.
- 2) Contact Medical Control to administer Midazolam/Diazepam for severe pain/muscle spasm.
- 3) Careful observation of respiratory status.

## PEDIATRIC ENVENOMATION – SNAKE BITES



- 1) Attempts to kill or capture the snake or bring dead animal to ED are NOT recommended.  
2) Many exotic snakes are neurotoxic so respiratory status must be monitored carefully.


**PEDIATRIC AIRWAY (1)**  
Airway Compromise  
I-99 Guideline see Appendix K



1) Medical Control contact is not mandatory, however, the medic is encouraged to discuss the situation with Medical Control if he/she is anticipating a Cricothyroidotomy and the clinical situation is such that there is time for Medical Control contact.  
 2) Verify proper tube placement by visualization of the cords and the tube passing through, bulb tube check/air aspiration, technique > 5 years old or EtCO2 detector/monitor for all ages, chest wall rise, good breath sounds, absence of gastric sounds, and clinical improvement in patient. Surgical Cricothyroidotomy contraindicated in children < 8 years old.  
 3)OG/NG tube placement if child ventilated with BVM for greater than 2 minutes or obvious gastric distention. Patients with head injuries should only have OG tube insertion, NG tube insertion contraindicated.

# PEDIATRIC AIRWAY OBSTRUCTED


I-99 Guideline see Appendix K



1) Verify proper tube placement by bulb tube check / air aspiration (if patient > 5 years old) or EtCO2 detector/monitor for all ages, chest wall rise, good breath sounds, absence of gastric sounds, and clinical improvement in patient. Surgical Cricothyroidotomy is contraindicated in patients < 8 years old.

## PEDIATRIC INSUFFICIENCY – BRONCHOSPASM

Applies to patients presenting with S/S of acute respiratory distress secondary to pre-existing condition or acute illness




- 1) Administer O<sub>2</sub> at high flow rates to all patients in severe respiratory distress. This is especially true if pulse oximetry is not available.
- 2) Consider the use of SVN therapy via in line BVM system in patients who are tiring or are appearing to have decreased tidal volumes.
- 3) If patients weight is less than 10 Kg, reduce Atrovent (Ipratropium Bromide) dose to 0.25 mg in 1.25 mL NS (½ unit dose)
- 4) Consider Epinephrine use in patients with poor tidal volumes or poor response to SVN.
- 5) Atrovent (Ipratropium Bromide) is contraindicated with soy or nut allergy.

# PEDIATRIC RESPIRATORY – UPPER AIRWAY EMERGENCIES

CROUP / EPIGLOTTIS


I-99 Guideline see Appendix K



- 1) BVM with reservoir with 100% O2 is usually adequate to provide ventilation and oxygenation. If ventilation appears clinically inadequate or transport will be greater than 5 minutes, consider intubation.
- 2) In patients 8 years old and under only needle cricothyrotomy with transtracheal jet insufflation should be utilized.
- 3) Medical Control contact is not mandatory, however, the medic is encouraged to discuss the situation with Medical control if he/she is anticipating a Cricothyroidotomy and the clinical situation is such that there is time for Medical Control contact.

## PEDIATRIC ALTERED LEVEL OF CONSCIOUSNESS

Altered level of consciousness and unconscious patient; includes GCS of 14 or less, psychotic or combative behavior, and the post seizure patient.




1) If opiate OD suspected BLS, management may be sufficient until response to Narcan (Naloxone) is determined.  
 2) Administer 0.5 – 1 Gm/kg of Dextrose. For neonates administer D<sub>10</sub> 2 mL/kg. For children less than one year of age administer D<sub>10</sub> 5-10mL/kg. For children 1-8 years of age, use D<sub>25</sub> 2-4 mL/kg. If unable to establish IV, give Glucagon 0.5 mg IM.  
 3) Infants and children < 20 kg or < 5 years receive 0.1mg/kg. Caution must be used in administration after birth to infants of addicted mothers, since it may precipitate abrupt narcotic withdrawal and seizures. Children older than 5 year or > 20 kg may be given up to 2.0 mg. Doses may be repeated at 2 minute intervals until narcotic reversal is achieved.  
 4) Possible causes/treatment could include: hypoxemia or acidosis (ventilate): hypovolemia (fluid bolus 20 mL/kg, repeat prn): tension pneumothorax (needle decompression): hyperthermia (cool patient): hypothermia (warm patient, monitor temperature): OD (examine scene): hypo/hyperglycemia (check blood glucose): postictal state (HPI/PMH)

# PEDIATRIC – SEIZURES OF UNKNOWN ETIOLOGY


Prolonged, Repetitive, or Status Epilepticus

I-99 Guideline see Appendix K




- 1) Administer 0.5 – 1 Gm/kg of Dextrose. For Neonates to one month old administer D10 2mL/kg. For children one month to one year of age administer D10 5-10mL/kg. For children 1-8 years of age, use D25 2-4 mL/kg. If unable to establish IV, give Glucagon 0.5 mg IM.  
 2) Do not administer Versed via the IN route if the patient's nose is bleeding or if nasal congestion or nasal discharge is present.  
 3) The only approved intranasal method of administering medications is with the Mucosal Atomizing Device (MAD)

## PEDIATRIC NAUSEA AND VOMITING




# PEDIATRIC ABDOMINAL PAIN, NON TRAUMATIC

I-99 Guideline see Appendix K




## PEDIATRIC ENVIRONMENTAL – HEAT RELATED




- 1) Do not cool below 102 degrees F.  
 2) Do not over cool and cause shivering and reoccurring heat buildup. If patient is shivering contact Medical Control to administer Midazolam or Diazepam.  
 3) If patient is agitated contact Medical Control to administer Midazolam or Diazepam.

## PEDIATRIC ENVIRONMENTAL - HYPOTHERMIA



- 1) If there is an organized rhythm do not begin CPR unless directed by Medical Control.  
 2) **Utilize only 1 shock.**  
 3) Contact Medical Control for ACLS medication administration regimen. Consider withholding medications if core temperature is  $\leq 86$  degrees F and an extended time between doses if temperature is  $> 86$  degrees F.


**PEDIATRIC  
HYPOTENSION / SHOCK, NON – TRAUMATIC  
I-99 Guideline see Appendix K**



- 1) BVM with reservoir with 100% O<sub>2</sub> and cricoid pressure is usually adequate to provide ventilation and oxygenation. If ventilation appears clinically inadequate or transport will be greater than 5 minutes, consider intubation.
- 2) If airway managed with BVM for > 2 minutes, insert 10-16 Fr OG/NG tube. Gastric decompression allows adequate pulmonary tidal volumes.
- 3) Rapid transport is of the utmost importance. Advanced life support procedures should be attempted at the scene, but if unsuccessful with a short period of time, the patient should be transported to the nearest appropriate facility without further delay.
- 4) Repeat assessment and lung auscultation before and after each fluid bolus.
- 5) If patient continues to be hypotensive, contact Medical Control to administer Dopamine 5-20 mcg/kg/min and/or Epinephrine infusion 0.1-1 mcg/kg/min.
- 6) Assess patient and patient symptoms to suggest cause and treat cause.

## PEDIATRIC POISONING / OVERDOSE (1)

I-99 Guideline see Appendix K



- 1) Patients who are suspected or known to have ingested substances with a suicidal intent may not refuse transport.
- 2) Administer 0.5 – 1 Gm/kg of Dextrose. For neonates administer D 10 2 mL/kg. For children less than one year of age administer D 10 5 – 10 mL/kg. For children 1-8 years of age, use D 25 2-4 mL/kg. If unable to establish IV, give Glucagon 0.5 mg IM.
- 3) Bring bottles / containers if possible. INSPECT SCENE.
- 4) Consider Medical Control input for Sodium Bicarbonate 1-2 mEq/kg for TCA overdose, Calcium Chloride 0.2 mL/kg very slow for calcium channel blocker overdose, Atropine 0.05 mg/kg every 2-4 min. for organophosphate exposure.

**APPENDIX A**  
**PEDIATRIC/NEONATAL STANDARDS/PHARMACOLOGICAL MODALITIES**

**PEDIATRIC/NEONATAL VITALS**

<b><u>AGE</u></b>	<b><u>HEART RATE/MIN</u></b>	<b><u>RESPIRATORY RATE/MIN</u></b>
Newborn	120 (70-180)	30 (30-60)
1 - 2 Years	120 (80-180)	27 (26-34)
2 - 4 Years	110 (80-140)	24 (20-30)
4 - 8 Years	100 (80-120)	22 (18-26)
8 - 12 Years	90 (70-110)	22 (15-24)

**BLOOD PRESSURE**

(\* Never inflate over 200 mmHg.)

(\* A convenient formula is:  $2 \times \text{age in years} + 70 = \text{Systolic}$ )

**WEIGHT**

(\* A convenient formula is:  $8 + \{2 \times \text{age in years}\} = \text{Weight in kilograms}$ )

**ENDOTRACHEAL TUBE**

(\* A convenient formula is:  $\frac{16 + \text{age in years}}{4} = \text{ET tube size}$ )

4

**PEDIATRIC LEVELS FOR DEFIBRILLATION**

Defibrillation energy level (2 joules/kg, double if unsuccessful)

Cardioversion energy level (0.5-1 joule/kg)

**PEDIATRIC PHARMACOLOGICAL MODALITIES**

Establishment of a pediatric IV line is frequently difficult or non-feasible in the field situation. Consider IO if situation dictates.

Dosages shown below are only to provide a standard. Actual dosage ordered by the responsible physician may be different.

Pediatric Age Clarification: VVMC Base Hospital will define the age to begin utilizing adult treatment guidelines as 14 years. In the case of the patient in cardiopulmonary arrest when the age is not known the AHA recommendation of using the presence of secondary sex characteristics as the determining factor of when to use guidelines is acceptable.

**PEDIATRIC  
DRUG ADMINISTRATION & PRECAUTIONS**

<b>ADENOSINE</b>	0.1 mg-0.2 mg/kg Rapid IV Push followed immediately by 2-3 mL NS. Monitor rhythm.
<b>AMIODARONE</b>	<b>VF/Pulseless VT</b> 5 mg/kg IV/IO (max 300mg single dose over 20 minutes may repeat every 5 min x 2) <b>SVT</b> 5 mg/kg IV/IO (max 150 mg single dose) over 20 minutes may repeat every 20 minutes x2 Mix Amiodarone only with D5W, max. dose is 15 mg/kg/day
<b>AMIODARONE DRIP</b>	1 MG/MIN FOR 6 HOURS, THEN 0.5 MG/MIN FOR UP TO 18 HOURS, MAXIMUM DAILY DOSE IS 15 MG/KG/DAY. Mix 450 mg in 250 ml D5W (special polyolefin bag) and run at 33.3 ml/hr for 1mg/min or 16.7 ml/hr for 0.5 mg/mi
<b>ATROPINE</b>	0.02 mg/kg IV/IO may repeat after 5 min. Min.: 0.1 mg Max: 0.5 mg child 1 mg adolescent
<b>CALCIUM CHLORIDE</b>	20 mg/kg IV/IO slowly over 10 min.
<b>DEXTROSE 50%</b>	(Peds) 0.5-1 Gm/kg diluted to D25W (Neonates) 0.1-0.2 Gm/kg diluted to D10W Slow IV/IO
<b>DIAZEPAM</b>	0.1-0.2 mg/kg IV/IO 0.5 mg/kg rectal to a total of 20 mg
<b>DIPHENHYDRAMINE</b>	1 mg/kg IV/IO/IM
<b>DOPAMINE</b>	2-20 mcg/kg/min. IV/IO Titrate to effect
<b>EPINEPHRINE 1:1000</b> (anaphylaxis/bronchospasm)	0.01 mg/kg IM Max 0.3 mg IM/SC
<b>IV EPINEPHRINE 1:10,000</b> (anaphylaxis)	0.01 mg/kg IV/IO Max single 0.5 mg, Titrate drip to effect

<b>EPINEPHRINE 1:10000</b> (cardiac)	0.01 mg/kg IV/IO may repeat every 3-5 minutes
<b>EPINEPHRINE DRIP</b>	0.1-1 mcg/kg/min. IV/IO Titrate to effect. For major cardiac events. Mix 4 mg 1:1000 in 250 mL NS for a 16 mcg/mL concentration
<b>FUROSEMIDE</b>	1 mg/kg IV/IO, Push slowly.
<b>GLUCAGON</b>	0.5 mg IM
<b>LIDOCAINE</b>	1 mg/kg IV/IO
<b>LIDOCAINE DRIP</b>	20-50 mcg/kg/min IV/IO When using 2 Gm/500 mL premix the concentration is 4000 mcg/mL
<b>METHYLPREDNISOLONE</b>	1-2 mg/kg IV/IO
<b>MIDAZOLAM</b>	0.05-0.1 mg/kg IV/IO Slowly over > 2 min may repeat every 2 min to a total of 10 mg 0.2 mg/kg IM to a total of 10 mg/0.2 mg/kg IN to a total of 10 mg
<b>MORPHINE</b>	0.1 mg/kg IV/IO/IM max dose 5 mg
<b>NALOXONE</b>	0.1 mg/kg SC/IV/IO If >5 yrs old or >20 kg 2 mg
<b>SODIUM BICARBONATE</b>	1 mEq/kg IV/IO Always dilute with sterile water or D5W 1:1 for infants up to 3 mos. Give slowly
<b>SVN: ALBUTEROL/ IPRATROPIUM</b>	2.5 mg/3 mL NS 0.02% 0.5 mg/2.5 mL NS, if < 10 Kg give half dose May repeat as necessary
<b>IV SOLUTIONS:</b>	
<b>RINGERS LACTATE</b>	20 mL/kg IV/IO Requires medical control input DO NOT USE on diabetic acidosis or hypothermia.
<b>NORMAL SALINE</b>	20 mL/kg IV/IO

APPENDIX B  
ADULT PHARMACOLOGICAL MODALITIES

Drug dosages listed on this page are intended as a general guideline for the usual dosages used in most situations. Expect to find variations from these standards.

<b>ADENOSINE</b>	6 mg IV/IO Rapid Push with 20 mL NS flush, may repeat in 1-2 min @ 12 mg x two repeats
<b>ALBUTEROL SULFATE-SVN</b>	2.5 mg/3 mL NS Unit Dose may repeat as necessary
<b>AMIODARONE</b>	<b>VF/Pulseless VT</b> 300 mg IV/IO push over 30-60 seconds, may repeat in 3-5 minutes with 150 mg once. <b>Wide-complex Tachycardia, AFib, Aflutter, SVT</b> 150 mg IV over 10 minutes (mix in 50 ml bag of D5W) may repeat every 10 minutes.
<b>AMIODARONE DRIP</b>	1 mg/min for 6 hours, then 0.5 mg/min for up to 18 hours. Maximum daily dose is 2.2 Mix 450 mg in 250 ml D5W (special polyolefin bag) and run at 33.3 ml/hr for 1mg/min or 16.7 ml/hr for 0.5 mg/mi
<b>ASA, BABY 81 mg</b>	4 chewable
<b>ATROPINE</b> - bradycardia asystole organophosphate poisoning	0.5 mg IV/IO, repeat every 5 min. to max of 3 mg 1 mg IV/IO 2 mg IV/IO repeat every 2-3 min prn titrate to atropinization
<b>CALCIUM CHLORIDE</b>	20 mg/kg of 10% Solution IV/IO for hyperkalemia and Ca Channel Blocker OD
<b>CHARCOAL ACTIVATED</b>	50 Gms
<b>DEXTROSE 50%</b>	25 Gms IV/IO Slow push
<b>DIAZEPAM</b>	2-10 mg Slow IV/IO. Titrate to effect.
<b>DILTIAZEM</b>	0.25 mg/kg IV slowly over 2 minutes, may repeat at 0.35 mg/kg in 15 minutes.
<b>DIPHENHYDRAMINE</b>	25-50 mg Slow IV/IM
<b>DOPAMINE</b>	5-20 mcg/kg/min IV/IO Drip
<b>EPINEPHRINE 1:1000</b>	0.1-0.3 mg IM

<b>EPINEPHRINE DRIP</b>	4 mg of 1:1000 Sol/250 mL D5W (16 mcg/mL concentration) <b>Initial dose 1 mcg/min.</b> Titrate to effect.
<b>EPINEPHRINE 1:10,000</b>	1 mg IV/IO
<b>FUROSEMIDE</b>	20 mg-80 mg IV/IO Slowly
<b>GLUCAGON</b>	1 mg IM - effect in 15-20 min
<b>IPRATROPIUM-SVN</b>	0.5 mg/2.5 mL NS Unit Dose, use with albuterol in first SVN only
<b>LIDOCAINE</b>	1 mg/kg IV/IO- Repeat 0.5 mg/kg every 5-10 min up to 3 mg/kg Cardiopulmonary arrest- 1.5 mg/kg repeat 0.75 mg/kg every 3-5 min. to 3 mg/kg
<b>LIDOCAINE DRIP</b>	2-4 mg/min IV/IO Drip
<b>MAGNESIUM SULFATE</b>	1-2 Gms in 50-100 mL D5W IV/IO over 2 min. (VF/pulseless VT - Give IV Push) PIH- 5 GM bolus in 50-100 ml D5W over 15 minutes then 1 –4 Gm/hr continuous infusion, mix 5 Gm/100 mL NS.
<b>METHYLPREDNISOLONE</b>	125 mg IV/IO
<b>MIDAZOLAM</b>	2.5 mg IV/IO slowly up to 10 mg 0.2 mg/kg IM to a total of 10 mg 0.2 mg/kg IN to a total of 10 mg
<b>MORPHINE SULFATE</b>	2-6mg IV/IM Slow IV push Initial dose max 2mg May repeat in 5 minutes for pain Max dose 20mg
<b>NALOXONE</b>	0.8 mg- 2 mg SC/IV every 3 min PRN
<b>NITROGLYCERIN</b>	0.4 mg (1/150) SL every 5 min X 3 if Systolic B/P > 100
<b>ONDANSETRONE</b>	4 mg slow IVP may repeat x 1 in 10 minutes if indicated
<b>SODIUM BICARBONATE</b>	1 – 2 mEq/kg IV/IO for wide QRS in tricyclic antidepressants overdose and hyperkalemia.
<b>THIAMINE</b>	100 mg IV/IM

## APPENDIX C

### VVMC PREHOSPITAL STANDARD INFUSION MIXTURES

**Amiodarone-** Mix 450 mg in 250mL of D5W (special polyolefin bag), concentration 1.8 mg/mL, and run at 33.3 mg/hr for 1 mg/min or 16.7 mL/hr for 0.5 mg/min

**Dopamine-** 400 mg/ 250 mL D5W premix= 1600 mcg/kg/min

**Epinephrine-** mix 4 mg 1:1,000/ 250 mL NS or D5W= 16 mcg/ mL

**Lidocaine-** 2 Gm in 500 mL D5W premix= 4 mg/ mL, run 1- 4 mg/min 15 to 60 gtts/min for adults.

4000 mcg/ mL to determine pediatric dosing of 20 -50 mcg/kg/min

**Magnesium Sulfate-** (OB Use) Mix 5 Gm/ 100 mL NS or D5W, run at 1-4 Gm/hr (20-80 mL/hr)

**Magnesium Sulfate-** (Asthma) Mix 2Gm/ 100mL NS, run over 10-20 minutes

**APPENDIX D**  
**AUTHORIZED SUPPLY OF MEDICATION FOR DRUG BOXES**  
**INTERFACILITY TRANSPORT MEDICATION LIST**

AGENT	MINIMUM SUPPLY	VVMC DRUG BOX
Adenosine	30 mg	6 mg / 2ml (5)
Albuterol Sulfate	10 mg	0.08% (6)
Amiodarone (optional)	300 mg	900 mg
Antiemetics: (optional) Promethazine HCL Ondansetron HCL Prochlorperazine edisylate	25 mg 4 mg 10 mg	8 mg
Aspirin	324 mg	81 mg (16)
Atropine Sulfate	4 mg 8 mg multi dose	1 mg/10 cc (4) 8mg/20 ml (1)
Calcium Chloride	1 gram	2
Charcoal, Activated (without sorbital)	50 G	25 gms (2)
Dexamethasone (optional)	8 mg	NONE
Dextrose	50 g	25g/50 ml (2)
Diazepam (required) Diazepam Rectal Delivery Gel (optional)		10mg/2ml (2)
Diltiazam (optional)	25 mg	1
Diphenhydramine HCL	50 mg	2
Dopamine HCL	400 mg	1
Epinephrine HCL 1: 1,000 solution	2 mg Multi-dose	1 cc amp (2) 30 cc (1)
Epinephrine HCL 1: 10,000 solution	5 mg	6
Etomidate (optional)	40 mg	2
Furosemide	100 mg	40mg/ml (4)
Glucagon	2 mg	2
Glucose, oral (optional)	30 gm	NONE
Ipratropium Bromide 0.02 %	5 ml	2.5 ml ud (4)
Lidocaine HCL IV Lidocaine Premixed Infusion	300 mg 2 G	100mg/5ml (4) 4mg/ml (500ml) (1)
Lorazepam (optional)	8 mg	NONE
Magnesium Sulfate	5 g	5
Methylprednisolone Sodium Succinate	250 mg	2
Midazolam (optional)	10 mg	5mg/5ml (2) 5mg/1ml (4)
Morphine Sulfate	20 mg	10 mg/ml (2)
Nalmefene HCL (optional)	4 mg	NONE
Naloxone HCL	10 mg	2 mg (5)
Nitroglycerin Tablets or	1 bottle	1

Nitroglycerin Sublingual Spray	1 bottle	
Oxytocin (optional)	10 units	20 units
Ondansetron (optional)	8 mg	8 mg
Phenylephrine Nasal Spray 0.5 %	1 bottle	1
Sodium Bicarbonate 8.4 %	100 mEq	50meq/50 ml (2)
Succinylcholine	20 mg/ml	200 mg (2)
Thiamine HCL	100 mg	1
Vasopressin	20 units	NONE
Verapamil HCL	10 mg	NONE
Nitrous Oxide (optional)	1 setup	NONE
Syringes: 1 ml TB	2	
3 ml	4	
10 ml	4	
20 ml	1	
50-60 ml	2	
Filter Needles	3	
Intravenous Solutions		
Dextrose 5% 250 ml (Optional)		1
Lactated Ringers 1000 ml	2	4
Normal Saline 1000 ml	2	4
250 ml	1	3
50 ml	2	2
<b>EMT BASIC DRUG BOX</b>		
Aspirin	324 mg	81 mg (16)
Epi- Auto injector	2 Adult	2
	2 Pediatric	2

**INTERFACILITY TRANSPORT MEDICATION LIST**

<b>IV INFUSIONS</b>	<b>EMT-P</b>	<b>EMT-I(99) Footnote 3</b>	<b>INFUSION PUMP</b>
AMIODARONE	X		X
ANTIBIOTICS	X	X	
BLOOD	X		
CALCIUM CHLORIDE	X		X
COLLOIDS	X	X	X
CORTICOSTEROIDS	X	X	X
DEXAMETHASONE	X		X
DILTIAZEM	X		X
DIURETICS- OTHER THAN FUROSEMIDE OR BUMETANIDE	X	X	X
DOPAMINE HCl	X		X
ELECTROLYTES/CRYSTALLOIDS- COMMERCIAL PREPARATIONS	X	X	
EPINEPHRINE HCL	X	X	X
FOSPHENYTOIN Na or PHENYTOIN Na	X		X
GLUCAGON	X	X	X
GLYCOPROTEIN IIb/IIIa Inhibitors	X		X
HEPARIN Na	X		X
H2 BLOCKERS	X	X	X

IV INFUSIONS	EMT-P	EMT-I(99) Footnote 3	INFUSION PUMP
MAGNESIUM SULFATE	X		X
METHYPREDNISOLONE SODIUM SUCCINATE	X	X	X
MIDAZOLAM	X		X
MORPHINE SULFATE	X	X	X
NITROGLYCERIN IV SOLUTION	X		X
OXYTOCIN	X		
PHENOBARBITAL Na	X		X
POTASSIUM SALTS	X		X
PROCAINAMIDE	X		X
RACEMIC EPINEPHRINE <i>svn</i>	X		
THEOPHYLLINE	X		X
TOTAL PARENTERAL NUTRITION	X		X
VITAMINS	X	X	

Notes:

1. Only an EMT-P may monitor an intravenous infusion via a central line.
2. Appropriate levels of EMT personnel shall be educated in an approved curriculum (covering both IV pumps and the specific drugs named in Table 1) and approved by their base hospital medical director, before monitoring patients on the listed medications during inter-facility transports.
3. On approval of the medical director on an individual basis.
4. All other medications an individual is authorized to administer by the Administrative Medical Director may be transported but must be on an infusion pump.

**APPENDIX E**  
**SCORES AND SCALES**

**Glasgow Coma Scale- Adult**  
**Level of Consciousness (LOC):**

1. Eye opening:	
Spontaneously	4
To speech	3
To pain	2
Never	1
2. Best verbal response	
Oriented	5
Confused	4
Inappropriate	3
Garbled	2
None	1
3. Best motor response	
Obeys commands	6
Localizes pain	5
Withdrawal	4
Abnormal flexion	3
Extension	2
None	1

Total = 3-15 possible

**Modified (Pediatric) Glasgow Coma Scale**

<b>Infants</b>		<b>Children</b>
	<b>Eye Opening</b>	
Spontaneous	4	Spontaneous
To speech or sound	3	To Speech
To painful stimulus	2	To pain
None	1	None
	<b>Best Verbal Response</b>	
Coos, babbles, smiles	5	Cries appropriately, Orientated
Irritable cry but consolable	4	Confused
Cries/screams to pain	3	Inappropriate crying/ Grunts incomprehensible words
Grunts/groans to pain	2	None
None	1	
	<b>Best Motor Response</b>	
Spontaneous movement	6	Obeys commands
Localizes pain	5	Localizes pain
Withdrawal from pain	4	Withdrawal from pain
Flexion to pain (decorticate)	3	Flexion to pain
Extension to pain (decerebrate)	2	Extension to pain
None	1	None

## Think F.A.S.T.

*For evaluation of acute, non-comatose, non-traumatic neurological complaint.*

Provide pre-arrival notification to Verde Valley Medical Center of potential stroke patient.

Notify by patch that you have a **Stroke Alert** if FAST score is 1 or greater.

<b>FACE</b>	<b>ARMS</b>	<b>SPEECH</b>	<b>TIME</b>
<b>Ask them to smile</b>	<b>Ask them to raise both arms</b>	<b>Ask them to speak a simple phrase</b>	<b>Time is BRAIN</b>
<b>Does the face look uneven?</b>	<b>Does one arm drift down?</b>	<b>Does the speech sound strange?</b>	<b>Time of symptom onset?</b>
<b>Yes= 1 point</b>	<b>Yes= 1 point</b>	<b>Yes= 1 point</b>	

### Exclusion Criteria:

- 
- **Seizure**
- **Stroke symptoms completely resolved**
- **Syncope**
- **Loss of consciousness**

## THE APGAR SCORE

**Appearance** (skin color):

Body and extremities blue, pale	0
Body pink, extremities blue	1
Completely pink	2

**Pulse rate:**

Absent	0
Below 100 bpm	1
100 bpm or more	2

**Grimace:**

No response	0
Grimace	1
Cough, sneeze, cry	2

**Activity:**

Limp	0
Some flexion of extremities	1
Active motion	2

**Respiratory effort:**

Absent	0
Slow and irregular	1
Strong cry	2

**Total score:** \_\_\_\_\_

APGAR Score should be assessed at one minute of birth and then reassessed at five minutes.

**APPENDIX F**  
**Transport Guidelines**

**VVEMS Medical Direction Policy on Transport Destination**

When ambulances are requested for a transport to a healthcare facility from the community, a private residence, doctors' offices and/or nursing homes they are to be transported to the closest, most clinically-appropriate facility.

Specific examples would include: Acute cerebral vascular accident (CVA), psychiatric patients, cardiology patients, and multi-trauma patients have specific destinations.

In cases when transport times are roughly equivalent, then considerations should be made on the destination facility based on the receiving facility's patient load or capacity, medical direction preference, and/or patient preference. Patient preference alone may not be sufficient reason to justify transport to a facility farther away than the closest most clinically-appropriate facility.

The goals of all EMS transports are to ensure the highest quality and safest patient care is being delivered while using public resources wisely, i.e. to minimize diversion of limited transport resources away from the community for extended periods of time. This philosophy will serve both patient and physicians' goals with an understanding that patient safety is the most important of these goals.

There may occur that reasonable circumstances in which a patient is best served by transport to a facility other than the closest. State EMS laws allow for these transports, but such transports shall occur subject to both online and administrative medical direction to govern these transport variances.

Under those limited circumstances in which patients may be transported to a facility other than the closest, the following criteria must be met:

1. Patient has been given informed consent to transfer and is aware that they are going to a facility farther away than the closest most appropriate facility.
2. The online medical direction physician (may be via nurse intermediary) has consented to the transport
3. If the transport is from a healthcare facility, both the sending and receiving physicians have consented to the transport and informed the patient of the risk/benefits of the transport to include most appropriate mode, i.e. ground vs. air transport.
4. The EMS crews on scene have determined the patient has been stabilized and is safe for transport to the more distant facility.
5. The EMS agency making the transport has sufficient personnel and resources to initiate the transport without delay or taking community transport resources without the ability to "backfill" the ambulance being sent on the transport.

**If any of the above criteria is not met, then the patient should be taken to the closest appropriate facility.**

**VVMC PREHOSPITAL CARE  
SEDONA FIRE DEPARTMENT TRANSPORT GUIDELINES**

**Sedona Emergency Center (SEC):**

SEC was established to provide rapid treatment of medical emergencies for the patients in the Sedona area. Patients should be transported to SEC unless their Chief Complaint falls into the categories outlined below.

All patients who need emergent stabilization will be transported to the closest facility

**Transport to VVMC:**

Cardiac: STEMI (transmit ECG ASAP).  
PTs from VOC with suspected cardiac in origin chest pain with or without STEMI.

Respiratory: Patients on CPAP mask who have improved and don't require immediate stabilization

Neurological: CVA—patients with acute onset of symptoms clearly less than 3 hours old

Psychiatric: Acute psychosis  
Acute suicidal ideation  
Combative and/or agitated\*\*

\*\*Any patients requiring security or constant nursing supervision for behavioral reasons are not appropriate for SEC

**Transport to FMC:**

Trauma: Any trauma patient who meets criteria for transport to a Trauma Center (see trauma transport guidelines)

Head Injury: Open skull fractures  
Significant facial injury  
Closed head injury with GCS less than 13 or any focal symptoms

Pediatrics: Patients who present with signs/symptoms of serious injury or illness (ex near drowning)

**These guidelines are not all inclusive. Final transport decision is at the discretion of the VVMC Base Station intermediary or physician.**

**APPENDIX G**  
RSI, CPAP, IO ACCESS

**RAPID SEQUENCE INTUBATION (RSI) USE BY EMT-P'S**

VVEMS medical direction supports the use of RSI as an optional advanced airway management skill by properly trained EMT-Ps in recognition of its potentially lifesaving results.

Purpose:

This airway management skill will be used in situations where placement of a prehospital endotracheal tube using RSI is indicated by patient conditions *and* where there is clear benefit of performing RSI in the prehospital environment.

Procedure:

1. EMT-P will work full-time for an agency that supports the optional RSI program.
2. EMT-P will complete the VVMC RSI training program before beginning to perform RSI in the field.
3. EMT-P will perform RSI using the Arizona Department of Health Services Recommendations for RSI in the field (ADHS, 2005).
4. EMT-P will complete a minimum of 12 patient or mannequin intubations a year to continue to be included in the RSI program.
5. EMT-P will complete an annual RSI refresher course.
6. EMT-P will participate in mandatory immediate self assessment and ongoing departmental CQI on all RSI in the field cases.
7. If requested, EMT-P will participate in review of cases through the Prehospital Peer Review Committee.

## CONTINUOUS POSITIVE AIRWAY PRESSURE (CPAP)

### PURPOSE:

Continuous Positive Airway Pressure has been shown to rapidly improve vital signs, gas exchange, and work of breathing, decrease the sense of dyspnea, and decrease the need for endotracheal intubation in patients who suffer from shortness of breath from asthma, COPD, pulmonary edema, CHF, and pneumonia. In patients with CHF, CPAP improves hemodynamics by reducing cardiac preload and afterload. CPAP decreases mortality when used in COPD exacerbations.

### INDICATIONS:

Any patient who is in respiratory distress of any cause who has protective airway reflexes AND

1. Is awake, oriented and able to follow commands
2. Is over 12 years old and is able to fit the CPAP mask
3. Has the ability to maintain an open airway
4. A respiratory rate greater than 25 breaths per minute
5. Has pulse oximetry less than 92%
6. Uses accessory muscles during respirations

### CONTRAINDICATIONS:

1. Patient is in respiratory arrest/apneic.
2. Patient is suspected of having a pneumothorax or has suffered trauma to the chest.
3. Patient has a tracheostomy.
4. Patient is actively vomiting or has upper GI bleeding.

### PRECAUTIONS:

1. Use care if patient:
  - a. Has impaired mental status and is not able to cooperate with the procedure
  - b. Has failed at past attempts at noninvasive ventilation
  - c. Complains of nausea or vomiting
  - d. Has inadequate respiratory effort
  - e. Has excessive secretions
  - f. Has a facial deformity that prevents the use of CPAP
2. Intubation should be performed by IEMT or Paramedic personnel if the patient:
  - a. Goes into respiratory or cardiac arrest
  - b. Is unresponsive to verbal stimuli (GCS is <9)
3. CPAP should not be used primarily with portable oxygen tanks because of the large amount of oxygen it takes to operate the device

### ADULT PROCEDURE:

1. Make sure patient does not have a pneumothorax!
2. Explain the procedure to the patient
3. Ensure adequate oxygen supply to ventilation device (100% when starting therapy and until SaO<sub>2</sub> is >92%)
4. Place the patient on continuous pulse oximetry
5. Place the patient on continuous endtidal CO<sub>2</sub> monitoring
6. Place patient on cardiac monitor and record rhythm strips with vital signs (interpretation by ALS personnel only)

7. Place the delivery device over the mouth and nose
8. Secure the mask with provided straps or other provided devices
9. Start CPAP at 5 cm H<sub>2</sub>O of PEEP. Increase gradually, if necessary, as patient adjusts and tolerates the PEEP to a maximum of 10 cm H<sub>2</sub>O on the pressure gauge. Document changes in patient status.
10. Check for air leaks
11. Monitor and document the patient's respiratory response to treatment
12. Check and document vital signs (ideally every 5 minutes)—specifically monitor rate, depth and SaO<sub>2</sub> and mental status. Some decrease in blood pressure may occur.
13. Continue to coach patient to keep mask in place and readjust as needed
14. Administer appropriate medication if necessary. (Ex. Albuterol/atrovent/methylprednisolone for asthma/COPD and Nitro for CHF)
15. If respiratory status deteriorates, remove device and consider intermittent positive pressure ventilation with or without endotracheal intubation
16. Contact receiving hospital in advance to advise them you have CPAP on the patient so they may prepare since equipment is not based in the ED.

#### REMOVAL PRODECURE:

1. CPAP therapy needs to be continuous and should not be removed unless the patient can not tolerate the mask or experiences continued or worsening respiratory failure or begins to vomit.
2. Intermittent positive pressure ventilation and/or intubation should be considered if the patient is removed from CPAP therapy.

#### PEDIATRIC CONSIDERATIONS:

1. CPAP should not be used in children under 12 years of age.

#### SPECIAL NOTES:

1. May be performed by Paramedics
2. May use .5 – 3 mg slow IV push of Midazolam if patient has high anxiety associated with CPAP device. Use small, repeated doses every 5 minutes.
2. Advise receiving hospital so they can be prepared for the patient
3. Do not remove CPAP until hospital therapy is ready to be placed on patient or if patient can no longer tolerate CPAP
4. Most patients will improve in 5-30 minutes. If no improvement within this time, consider intermittent positive pressure ventilation
5. Watch patient for gastric distention
6. Be cautious when using nitroglycerine spray with CPAP since it could be dispersed on EMT's

## EMERGENT RESPIRATORY CPAP DEVICE ONLY

### SETTING THE CPAP LEVEL

Prior to setting the pressure always observe that the airway pressure gauge needle indicator is at the zero (0) value with the CPAP adjustment knob in the fully counterclockwise position and the breathing circuit connected. The zero point may be adjusted by adjusting the small screw on the face of the gauge. To set continuous positive airway pressure, turn the CPAP adjustment clockwise and observe the needle indicator on the airway pressure gauge.

### APPLYING THE BREATHING CIRCUIT AND MASK

Assembly Instructions:

The CPAP Breathing Circuit is pre-assembled at the factory. To attach the breathing circuit to the CPAP unit insert and align the locking bayonet outlet adapter to the unit and turn clockwise until securely engaged.

1. When the mask is ready and the patient is prepared, turn the ON/OFF valve fully to the ON position (counterclockwise ½ turn). Ensure that the gas is flowing, and then hold the mask to the patient's face. Gently place your other hand on the back of the patient's head to confirm a good air seal.
2. Within a few minutes (once the patient is comfortable) use the head strap to hold the mask in place. Ensure that the mask is not too tight. Some air leakage is acceptable unless it is in the eye area.
3. Make sure the flow rate is in excess of the patient's inspiratory flow rate in order to maintain continuous pressure throughout the breathing cycle. Check this frequently during transport as the patient's needs may change. There are 3 ways to determine whether the flow is set high enough:
  - a. The CPAP valve should remain slightly open during the entire respiratory cycle.
  - b. The anti-asphyxia valve on the mask should not open during normal operation.
  - c. Some gas should escape from the exhalation port.
4. For patient comfort, and to preserve oxygen, turn the flow adjustment knob down to maintain the flow just above the patient's flow rate.
5. In most cases, the patient should improve in the first 5 minutes with CPAP.
6. If after 5 minutes the patient's SpO<sub>2</sub> is not at the desired level, deliver higher oxygen concentrations (up to 100%) by turning the valve farther counterclockwise. The FiO<sub>2</sub> should be increased judiciously to preserve O<sub>2</sub>. Adjust one to two full turns and then re-evaluate the SpO<sub>2</sub>.

### DOCUMENTATION FOR ADHERENCE TO PROTOCOL:

The following items must be documents for Adherence to Protocol:

1. Prehospital impression as to why CPAP was chosen
2. Vital signs (BP, HR, RR, SpO<sub>2</sub>) recorded every 5 minutes
3. Description of patient's response to CPAP
4. Documentation of other airway adjunct if CPAP is unsuccessful
5. Use of sedating medications

## IO Protocol for use with Easy IO Gun

### **Training:**

EZ-IO® infusion systems require specific training prior to use.

### **INDICATIONS:**

EZ-IOAD (40 kg and over) EZ-IO® PD (3 – 39 kg) and EZ-IO LD

Note: *Certain patients may require a needle set outside their ideal weight range “One size needle set does not fit all”*

1. Immediate vascular access in emergencies.
2. Intravenous fluids or medications are urgently needed and a peripheral IV cannot be established in 2 attempts or 90 seconds

**AND** the patient exhibits risk of immediate death or loss of function or deterioration.

### **Relative CONTRAINDICATIONS:**

Fracture of the bone selected for IO infusion (*consider alternate sites*)

Excessive tissue at insertion site with the absence of anatomical landmarks (*consider alternate sites*)

Previous significant orthopedic procedures (*IO within 24 hours, prosthesis - consider alternate sites*)

Infection at the site selected for insertion (*consider alternate sites*)

### **CONSIDERATIONS:**

**Flow rate:** Due to the anatomy of the IO space, flow rates may appear to be slower than those achieved with an IV catheter.

- Ensure the administration of an appropriate rapid **SYRINGE BOLUS (flush)** prior to infusion  
“**NO FLUSH = NO FLOW**”
  - Rapid syringe bolus (flush) the EZ-IO AD or LD with 10 ml of normal saline
  - Rapid syringe bolus (flush) the EZ-IO PD with 5 ml of normal saline
  - Repeat syringe bolus (flush) as needed
- To improve continuous infusion flow rates always use a syringe, pressure bag or infusion pump

**Optional treatment for Pain after stabilization of patient:** IO Infusion for conscious patients has been noted to cause severe discomfort

- **SLOWLY** administer Lidocaine 2% (Preservative Free) through the EZ-IO hub. *Ensure that the patient has no allergies or sensitivity to Lidocaine.*
  - EZ-IO AD and EZ-IO LD Slowly administer 20 – 40 mg Lidocaine 2% (Preservative Free)
  - EZ-IO® PD Slowly administer 0.5 mg /kg Lidocaine 2% (Preservative Free)

### **EQUIPMENT:**

EZ-IO Driver

EZ-IO AD, EZ-IO PD or EZ- IO LD Needle Set

Alcohol or Betadine Swab

EZ-Connect® or Standard Extension Set

10 ml Syringe

Normal Saline (or suitable sterile fluid)

Pressure Bag or Infusion Pump  
2 % Lidocaine (preservative free)  
EZ-IO® Yellow wristband

**PROCEDURE:** *If the patient is conscious, advise of EMERGENT NEED for this procedure and why*

1. Wear approved Body Substance Isolation Equipment (BSI) or Personal Protective Equipment (PPE)
2. Determine EZ-IO® Indications
3. Rule out Contraindications
4. Locate appropriate insertion site (Preferred sites: *Proximal / Distal Tibia*  
*Approved site: Proximal Humerus*)
5. Prepare insertion site using aseptic technique
6. Prepare the EZ-IO® driver and appropriate needle set
8. Stabilize site and insert appropriate needle set
9. Remove EZ-IO® driver from needle set while stabilizing catheter hub
10. Remove stylet from catheter, place stylet in shuttle or approved sharps container
11. Confirm placement
12. Connect primed EZ-Connect®
13. Slowly administer appropriate dose of Lidocaine 2% (Preservative Free) IO to conscious patients
14. Syringe bolus (flush) the EZ-IO® catheter with the appropriate amount of normal saline.
15. Begin infusion with pressure (syringe bolus, pressure bag or infusion pump)
17. Dress site, secure tubing and apply wristband as directed
18. Monitor EZ-IO® site and patient condition – Remove catheter within 24 hours.

**APPENDIX H**  
**HEAD TO TOE ASSESSMENT**

**The Head-to Toe Assessment should include these areas:**

1. Complete set of vital signs
2. Head:
  - a) Inspect and palpate scalp, face, ears, nose, eyes
  - b) Check pupils for size, equality, reaction to light, accommodation
3. Neck:
  - a) Inspect and palpate location of trachea
  - b) Check jugular veins
  - c) Palpate cervical spine
4. Chest/Back:
  - a) Inspect, palpate, and auscultate chest and back
5. Abdomen/Pelvis/Buttocks:
  - a) Inspect, palpate, auscultate abdomen
  - b) Perform 3 point pelvis check
6. Lower Extremities
  - a) Inspect and palpate both legs and feet
  - b) Check circulation, sensation, and motor function in both feet
7. Upper Extremities:
  - a) Inspect and palpate both arms and hands
  - b) Check circulation, sensation, and motor function in both hands
8. Neurological Assessment
9. EKG Monitoring/12 Lead
10. Pulse Oximetry
11. Glucose Determination
12. History

**APPENDIX I**  
**ALS Release of Patients for BLS Transport**

**Criteria 1: Non-emergency category must have vitals within the following limits:**

Adult

- \*Respirations 10 to 24
- \*BP 90 to 160 systolic  
60 to 110 diastolic
- \*Pulse 60 to 100
- \*Pulse Oximetry >90% or change from normal

Pediatric

Age Appropriate



**Criteria 2: The following high-risk indications must be absent:**

- Abdominal pain- Adult
- Altered mental status (Compared to pt's normal status)
- Any acute cardiac arrhythmia
- Chest pain
- Shortness of breath
- Syncope/ Dizziness
- Overdose/poisoning
- Seizures
- Pregnancy- related complaint
- Significant head/neck/chest/abdomen/pelvis trauma



**Criteria 3: Absence of disease or process that would benefit from ALS care**

A physical exam must be completed and documented. After evaluation the patient must not have any signs or symptoms that would indicate significant findings or emergent condition. Patient care may be upgraded to ALS at anytime if medic feels patient warrants additional care.



Contact must be made to medical control for final approval to transport BLS. BLS provider may complete courtesy notification with the guidance of ALS provider.

## VVMC IEMT Care Levels

<b>Skills- Airway/Ventilation/Oxygenation</b>	<b>Yes=IEMT may do. No EMT May not do.</b>
Airway-esophageal	<b>Yes</b>
Airway-supraglottic	<b>Yes</b>
Airway-nasal	<b>Yes</b>
Airway-oral	<b>Yes</b>
Bag-valve-mask (BVM)	<b>Yes</b>
BiPAP/CPAP	<b>Yes</b>
Chest decompression-needle	<b>Yes</b>
Chest tube placement-assist only	<b>No</b>
Cricoid pressure (Sellick's maneuver)	<b>Yes</b>
Cricothyrotomy-needle	<b>No</b>
Cricothyrotomy-percutaneous	<b>No</b>
Cricothyrotomy-surgical	<b>No</b>
Demand valve-manually triggered ventilation	<b>Yes</b>
End tidal CO2 monitoring/capnography	<b>No</b>
Gastric decompression-NG tube	<b>Yes</b>
Gastric decompression- OG tube	<b>Yes</b>
Head Tilt chin lift	<b>Yes</b>
Intubation-nasotracheal	<b>No</b>
Intubation-oro-tracheal	<b>Yes</b>
Jaw-thrust	<b>Yes</b>
Jaw-thrust- modified (trauma)	<b>Yes</b>
Medication Assisted Intubation	<b>No</b>
Mouth-to-barrier	<b>Yes</b>
Mouth-to-mask	<b>Yes</b>
Mouth-to-mouth	<b>Yes</b>
Mouth-to-nose	<b>Yes</b>
Mouth-to-stoma	<b>Yes</b>
Obstruction-direct laryngoscopy	<b>Yes</b>
Obstruction-manual	<b>Yes</b>
Oxygen therapy-humidifiers	<b>Yes</b>
Oxygen therapy-nasal cannula	<b>Yes</b>
Oxygen therapy-non rebreather mask	<b>Yes</b>
Oxygen therapy-partial non-rebreather	<b>Yes</b>


<b>Skill-Airway Continued</b>	<b>Yes/No</b>
Oxygen therapy-simple face mask	<b>Yes</b>
Oxygen therapy-venturi mask	<b>Yes</b>
PEEP-therapeutic	<b>No</b>
Pulse Oximetry	<b>Yes</b>
Suctioning-upper airway	<b>Yes</b>
Suctioning-tracheobronchial	<b>Yes</b>
Automated transport ventilator	<b>Yes</b>
<b>Skill-Cardiovascular/Circulation</b>	<b>Yes/No</b>
Cardiac monitoring –multiple lead interpretive	<b>No</b>
Cardiac monitoring-single lead interpretive	<b>YES</b>
Cardiopulmonary resuscitation	<b>Yes</b>
Cardioversion-electrical	<b>No</b>
Carotid massage	<b>No</b>
Defibrillation-automatic/semiautomatic	<b>Yes</b>
Debrillation-manual	<b>Yes</b>
Hemorrhage control-direct pressure	<b>Yes</b>
Hemorrhage control-tourniquet	<b>Yes</b>
Internal: cardiac pacing-monitoring only	<b>Yes</b>
MAST/PASG	<b>Yes</b>
Mechanical CPR device	<b>Yes</b>
Transcutaneous pacing-manual	<b>Yes</b>
<b>Immobilization</b>	<b>Yes/No</b>
Spinal immobilization-cervical collar	<b>Yes</b>
Spinal immobilization-long board	<b>Yes</b>
Spinal immobilization-manual	<b>Yes</b>
Spinal immobilization-seated (KED)	<b>Yes</b>
Spinal immobilization-rapid manual extrication	<b>Yes</b>
Extremity stabilization-manual	<b>Yes</b>
Extremity splinting	<b>Yes</b>
Splint-traction	<b>Yes</b>
Mechanical patient restraint	<b>Yes</b>
Emergency moves for endangered patients	<b>Yes</b>

<b>Medication administration-routes</b>	<b>Yes/No</b>
Assisting patient with his/her own prescribed medications (aerosolized/nebulized)	<b>Yes</b>
Aerosolized/nebulized (beta agonist)	<b>Yes</b>
Buccal	<b>Yes</b>
Endotracheal tube	<b>Yes</b>
Inhaled self-administered (nitrous oxide)	<b>Yes</b>
Intradermal	<b>No</b>
Intramuscular	<b>Yes</b>
Intranasal	<b>Yes</b>
Intravenous push	<b>Yes</b>
Intravenous piggyback	<b>Yes</b>
Nasogastric	<b>No</b>
Oral	<b>Yes</b>
Rectal	<b>Yes</b>
Subcutaneous	<b>Yes</b>
Sublingual	<b>Yes</b>
Auto-injector (self or peer)	<b>Yes</b>
Auto-injector (patients own prescribed medications)	<b>Yes</b>
<b>IV initiation/maintenance fluids</b>	<b>Yes/No</b>
Access indwelling catheters and implanted central IV ports	<b>No</b>
Central line-monitoring	<b>No</b>
Intraosseous-initiation	<b>Yes</b>
Intravenous access	<b>Yes</b>
Intravenous initiation-Central	<b>No</b>
Intravenous initiation-peripheral	<b>Yes</b>
Intravenous-maintenance of non-medicated IV fluids	<b>Yes</b>
Umbilical initiation	<b>No</b>
<b>Miscellaneous</b>	<b>Yes/No</b>
Assisted delivery (childbirth)	<b>Yes</b>
Assisted complicated delivery (childbirth)	<b>Yes</b>
Blood glucose monitoring	<b>Yes</b>
Blood pressure-automated	<b>Yes</b>

Blood pressure-manual	<b>Yes</b>
Eye irrigation	<b>Yes</b>
Eye irrigation (Morgan lens)	<b>No</b>
Thrombolytic therapy-initiation	<b>No</b>
Urinary catheterization	<b>No</b>
Venous blood sampling	<b>Yes</b>
Blood chemistry analysis	<b>No</b>
Interfacility med transport list	<b>See next section</b>
<b>Medications</b>	<b>Yes/No</b>
Adenosine	<b>Yes</b>
Albuterol Sulfate (SVN or MDI)	<b>Yes</b>
Amiodarone	<b>Yes</b>
Antibiotics	<b>Transport Agent</b>
Aspirin	<b>Yes</b>
Atropine Sulfate	<b>Yes</b>
Apropine Sulfate (multidose vial)	<b>Yes</b>
Atropine Sulfate Auto-injector	<b>Yes</b>
Atropine Sulfate and Pralidoxime Chloride (combined)	<b>Only in Toxicological Emergencies</b>
Blood	<b>No</b>
Bronchodilator, inhaler	<b>Patient Assist Only</b>
Calcium Chloride	<b>No</b>
Calcium Gluconate, 2.5% topical gel	<b>Topical Agent- Only in Toxicological Emergencies</b>
Charcoal, Activated	<b>Yes</b>
Colloids	<b>Transport Agent</b>
Corticosteroids	<b>Transport Agent</b>
Dexamethasone	<b>Yes</b>
Dextrose D50	<b>Yes</b>
Dextrose, 5% in H2O	<b>Yes</b>
Diazepam	<b>Yes</b>
Diazepam Rectal Delivery Gel	<b>Yes</b>
Diltiazem or Verapamil HCl	<b>No</b>
Diphenhydramine HCl	<b>Yes</b>
Diuretics	<b>Transport Agent</b>
Dopamine HCl	<b>No</b>
Electrolytes/Crystalloids (commercial preparations)	<b>Transport Agent</b>
Epinephrine Auto-injector	<b>Yes</b>
Epinephrine HCl, 1:1000	<b>Yes</b>
Epinephrine HCl, 1:1000 Multidose	<b>Yes</b>
Epinephrine HCl, 1:10000	<b>Yes</b>
Etomidate	<b>No</b>

Fosphenytoin Na or Phenytoin Na	<b>No</b>
Furosemide or Bumetanide	<b>Yes</b>
Glucagon	<b>Yes</b>
Glucose, Oral	<b>Yes</b>
Glycoprotien IIb, IIIa Inhibitors	<b>No</b>
H2 Blockers	<b>Transport Agent</b>
Heparin Na	<b>No</b>
Immunizing Agent	<b>Yes</b>
Ipratropium Bromide 0.02%	<b>Yes</b>
Lactated Ringers	<b>Yes</b>
Lidocaine HCl IV	<b>Yes</b>
Lorazepam	<b>Yes</b>
Magnesium Sulfate	<b>No</b>
Methylprednisolone Sodium Succinate	<b>Yes</b>
Midazolam	<b>No</b>
Morphine Sulfate	<b>Yes</b>
Nalmefine HCl	<b>Yes</b>
Naloxone HCl	<b>Yes</b>
Nitroglycerin IV Solution	<b>No</b>
Nitroglycerin Sublingual Spray or Nitroglycerin Tablets	<b>Yes</b>
Nitrous Oxide	<b>Yes</b>
Normal Saline	<b>Yes</b>
Ondansetron HCl	<b>Yes</b>
Oxygen	<b>Yes</b>
Oxytocin	<b>Yes</b>
Phenobarbital Na	<b>No</b>
Phenylephrine Nasal Spray	<b>Yes</b>
Potassium Salts	<b>No</b>
Pralidoxine Chloride, Auto-ejector	<b>Only in Toxicological Emergencies</b>
Procainamide HCl	<b>No</b>
Racemic Epinephrine	<b>No</b>
Sodium Bicarbonate 8.4%	<b>Yes</b>
Succinylcholine	<b>No</b>
Theophylline	<b>No</b>
Thiamine HCl	<b>Yes</b>
Total Parenteral Nutrition, with or without lipids	<b>No</b>
Tuberculin PPD	<b>Yes</b>
Vasopressin	<b>No</b>
Vitamins	<b>Transport Agent</b>


## Appendix K WITHHOLDING/TERMINATION OF RESUSCITATION



(1) For indications where no treatment is required the provider may withhold resuscitative efforts and patch for medical direction.  
(2) Consideration should be given to potentially reversible conditions such as, overdose/poisoning, hypothermia, cold water drowning, etc.

## ADULT BRADYCARDIA, UNSTABLE (1)


HEART RATE < 60 MINUTE WITH ACCOMPANYING SIGNS/SYMPTOMS OF HEMODYNAMIC COMPROMISE, I.E., CHEST PAIN, HYPOTENSION, IF HISTORY/EVIDENCE OF TRAUMA, PROCEED TO TRAUMA TREATMENT GUIDELINE



- (1) Signs/symptoms of an unstable patient may include chest pain, SOB, decreased LOC, hypotension, shock, pulmonary edema, congestive heart failure, and acute myocardial infarction.  
 (2) This should not delay definitive treatment.  
 (3) Repeat vital signs and lung auscultation before and after fluid administration


### CHEST PAIN SUGGESTIVE OF CARDIAC ORIGIN

Chest Pain suggestive of possible myocardial ischemia (1)




- 1) Indications of chest pain suggestive of possible myocardial ischemia include: Description of crushing, squeezing, pressure, burning, tightness, diaphoresis, nausea/vomiting, apprehension, radiation, age>30, **associated cardiac risk factors.**
- 2) If 3 lead capability, should be done in pt's initial assessment.
- 3) Repeat vital signs and lung auscultation before and after administration of NTG. Consider prior NTG use. If pain reoccurs and is not refractory to NTG, repeat NTG 0.4mg SL every 5 minutes as needed for pain relief, maintaining B/P > 100.
- 4) Nitroglycerin is contraindicated in patients that have taken Viagra (sildenafil), Cialis, Levitra, or similar medications in the previous 72 hours.
- 6) Communication with hospital should be completed as soon as possible so that Cath Lab team can be notified for ST Elevation MI.

**ADULT PULSELESS ARREST**




- 1) Assess rhythm – quick look, only check pulses if there is an organized rhythm present.
- 2) Evaluate airway, intubate if necessary, limit interruption of CPR as much as possible.
- 3) Once patient is successfully intubated perform continuous asynchronous compression (rate 100/min) with ventilations (rate 8-10/min)
- 4) Pulse checks should be done only if EKG indicates a potentially perfusing rhythm, do not interrupt chest compressions, and be very brief.
- 5) Medications should be administered during CPR as soon as possible after rhythm checks.
- 6) Consider possible causes: Hypovolemia,(volume infusion), hypoxia (ventilation/re-evaluation),acidosis (ventilation/re-evaluation), tension pneumothorax (needle decompression), hypothermia, hypoglycemia, drug overdose, cardiac tamponade (volume infusion), massive AMI, hyperkalemia (consider NaHCO<sub>3</sub>, D50W) massive pulmonary embolism.
- 7) If patient remains asystolic or other agonal rhythm after successful intubation, initial medications, no reversible causes are identified, and transport has not been initiated, consider termination of resuscitative efforts by order of a physician. Consider interval since arrest.
- 8) For successful conversions with HR>60 and no 2<sup>nd</sup> or 3<sup>rd</sup> degree heart blocks. Assess vital signs, administer Lidocaine 1-1.5 mg/kg and start infusion at 2-4 mg/ min. If patient received bolus doses prior to conversion administer maintenance infusion only.

## CARDIAC ARREST POST RESUSCITATION INDUCED HYPOTHERMIA




- 1) Criteria for Induced Hypothermia:
  - ROSC after cardiac arrest not related to trauma or hemorrhage
  - Age greater than 18 years old
  - Patient is intubated and remains comatose (no purposeful response to pain)
  - GCS of less than 8 after ROSC
  - ROSC within 60 minutes of arrest
  - Initial temperature >34C- no environmental hypothermia related arrhythmia
  - Female patients not pregnant
  - No uncontrolled hemorrhage
  - No persistent unstable arrhythmia
  - No DNR paperwork identified during resuscitation
- 2) If unable to intubate DO NOT initiate induced hypothermia
- 3) Patients develop metabolic alkalosis with cooling. Do not hyperventilate.
- 4) When exposing pts for cooling purposes undergarments may remain in place. Be mindful of your environment and be preserve patients modesty.
- 5) Do not delay transport for purposes of cooling.
- 6) Patch early to advise of induction of hypothermia to allow ED to have equipment ready.
- 7) **AT ANY TIME- Loss of Spontaneous Circulation- discontinue cooling and go to appropriate protocol**

## ADULT TACHYCARDIA WITH PULSES




- 1) If at any time patient becomes unstable, proceed to "Unstable" side
- 2) Contact medical control for Amiodarone 150 mg administered over 10 minutes, if no response may repeat every 10 minutes, maintenance infusion after conversion is 1 mg/min.
- 3) For successful conversions of ventricular arrhythmias with HR > 60 and no 2<sup>nd</sup> or 3<sup>rd</sup> degree heart blocks: Assess vital signs, administer Amiodarone 150 mg IV over 10 minutes then begin drip at 1 mg/min. for first 6 hours or Lidocaine (Xylocaine) 1mg/kg and start infusion at 2-4 mg/min, reduce maintenance infusion of Lidocaine by half in patients with renal or hepatic disease or > 70 years of age. If patient received bolus doses prior to conversion administer maintenance infusion only.
- 4) If rhythm is polymorphic VT go immediately to unsynchronized defibrillation at 120-200 biphasic with manufactures recommendations or monophasic 360J. For polymorphic VT the provider should be prepared to move immediately to the Pulseless Arrest algorithm if pulseless arrest develops.

### TRAUMA – SPINAL INJURY




## ANAPHYLAXIS

Applies to patient presenting with allergic reaction and with signs and symptoms of airway, respiratory, or circulatory compromise (laryngeal edema, bronchospasm, or hypotension. )



- 1) If signs and symptoms of severe hypoperfusion and an IV can be rapidly established, consider going directly to IV Epinephrine as per protocol.
- 2) Establishment of an IV should not delay the administration of IM Epinephrine to a patient in extremis.
- 3) The use of Epinephrine in patients age > 45 years or with known coronary artery disease requires Medical Control input.
- 4) If prolonged transport consider repeat use of Epinephrine every 15 minutes. Medical Control input should be obtained, if possible.
- 5) At any time an IV cannot be established, give Benadryl ( Diphenhydramine) 50 mg IM as soon as possible after Epinephrine IM.
- 6) Consider IO if no IV access and patient is in extremis.
- 7) Consider the use of SVN therapy via in line BVM system in patients who are tiring or are appearing to have decreased tidal volumes.
- 8) If patient continues to be hypotensive contact Medical Control to administer Epinephrine 1:1,000 2-10 mcg/min IV/IO Infusion, titrate to effect.


## CEREBRAL VASCULAR ACCIDENT - STROKE



1) Establishing time signs and symptoms began is CRITICAL. If patient awoke from sleep with S/S it is also important to determine how long patient was asleep. Patients with ischemic strokes < 3 hours old may be candidates for TPA therapy with some candidates eligible for up to 4.5 hours.


2) Do not treat patients for hypertension unless directed to do so by medical control.

## ENVENOMATION - ARACHNIDS



- (1) Attempts to kill or capture insect or bring to ED are not recommended.
- (2) Contact Medical Control to administer Valium (Diazepam) for severe pain / muscle spasm.
- (3) Careful observation of respiratory status.


### AIRWAY COMPROMISE



1) Consider OG tube for gastric decompression in situations of prolonged ventilation.  
 2) Consider blind airway insertion device if difficult airway and Endotracheal intubation unsuccessful after two attempts.


### RESPIRATORY INSUFFICIENCY – BRONCHOSPASM

Applies to patients with S/S of acute respiratory distress, secondary to asthma, COPD, and inhalation injury




- 1) Administer O2 at high flow rates to all patients in severe respiratory distress. This is especially true if pulse oximetry is not available.
- 2) Consider the use of SVN therapy via in line BVM system in patients who are tiring or are appearing to have decreased tidal volumes.
- 3) The use of epinephrine in patients 45 years or greater or with known coronary artery disease requires **Medical Control** input.
- 4) Do not delay definitive therapy to establish IV.
- 5) Obtain an IO if no IV access and patient is in extremis.
- 6) Atrovent (Ipratropium Bromide) is contraindicated in patients with soy or nut allergy.
- 7) Epinephrine IM is indicated for use in bronchospasm i.e. bronchiolitis and asthma

### AIRWAY OBSTRUCTED




1) Chest thrusts if patient is obese or pregnant.  
2) Verify proper tube placement by bulb tube check / air aspiration, chest wall rise, good breath sounds, absence of gastric sounds, and clinical improvement in patient.

## RESPIRATORY INSUFFICIENCY – PULMONARY EDEMA



- 1) Patients who appear to be tiring or have decreased tidal volume may require respiratory assist.
- 2) High flow O<sub>2</sub> should be used in any patient who appears distressed
- 3) Repeat vital signs and lung auscultation before and after administration of NTG.
- 4) Unless directed by online medical control Nitroglycerin is contraindicated in patients that have taken Viagra (sildenafil), Cialis, Levitra, or similar medication within 72 hours


## OBSTETRICS COMPLICATIONS OF PREGNANCY



1) Signs of PIH/ pre-eclampsia / eclampsia may include: Diastolic BP . 80 mmHg with cerebral or visual disturbances, epigastric or RUQ pain with nausea and vomiting, ALOC, hyper-reflexia, peripheral edema, pulmonary edema, seizures.  
 2) Contact Medical Control to administer 1 liter fluid bolus of NS and consideration for very mild sedation with diazepam

**ALTERED LEVEL OF CONSCIOUSNESS**


GCS of 14 or <, psychotic or combative behavior, the post seizure patient, the near/post syncope patient, or any patient with history of ALOC as a part of current event.(1)



- 1) Utilize information obtained from family, bystanders, friends, or other health care workers.
- 2) If hypoglycemia or opiate OD suspected, BLS airway management may be sufficient until response to Dextrose and/ or Naloxone is determined.
- 3) If no history of alcoholism is suspected and malnutrition or cachexia is not present, Thiamine may be withheld.
- 4) If no change in LOC, repeat glucose. Realize the onset of action of Glucagon is 5-15 minutes.


### ALTERED LEVEL OF CONSCIOUSNESS

With suspected alcohol intoxication (1)



- 1) Utilize information obtained from family, bystanders, friends, or other health care workers.
- 2) If hypoglycemia or opiate OD suspected, BLS airway management maybe sufficient until response to Dextrose and/ or Narcan ( Naloxone) is determined.
- 3) If no change in LOC, repeat glucose. Realize the onset of action of Glucagon is 5-15 minutes.
- 4) Do not intubate sleeping, stable intoxicated patient if oxygenating and ventilating.


## ADULT VIOLENT/AGITATED PATIENT (1)



- (1) Agitated patient (significant psychomotor agitation) due to possible drug ingestion and/or violent patients who after initial treatment remain a danger to self or others.
- (2) Patients may need to be restrained for patient, provider, or bystander safety in order to complete other appropriate treatment guidelines.
- (3) If patient is in police custody and handcuffs have been applied it is preferable that a police officer also accompany the patient. EMS providers must, at a minimum, have the handcuff key in their possession during transport.
- (4) Patients shall be positioned in a manner that does not compromise airway or breathing. No patient will be restrained prone or "hog-tied." No patient will be placed between backboards or gurneys.


### SEIZURE

Prolonged, Repetitive, or Status Epilepticus




## ABDOMINAL PAIN, NON – TRAUMATIC

Testicular torsion, Pelvic pain, AAA



1) PASG is a class 1 recommendation for hypovolemia due to ruptured abdominal aortic aneurysm.  
2) Start initial Morphine dose at 2mg to determine if patient tolerates narcotics.


**ENVIRONMENTAL – HEAT RELATED**



1) Do not cool below 102 degrees F.  
 2) Do not over cool and cause shivering and reoccurring heat buildup. If pt is shivering contact Medical Control to administer Diazepam.  
 3) If patient is agitated contact Medical Control to administer Diazepam.


### HYPOTENSION, NOT TRAUMATIC (1)

Applies ONLY when other specific ALS protocols do not apply.  
 Hypotension is defined as BP < 90 systolic and associated signs / symptoms of hypoperfusion.  
 If history / evidence of Trauma, proceed to Trauma Treatment Guideline.




- 1) PMH and patient's medications may be key to index of suspicion for cause of hypotension, e.g. history of ulcers, aneurysm, previous cardiac disease, alcoholism, etc. Consider possible causes of hypotension and treat cause.
- 2) Consider establishing 2 large bore IV's dependent upon patient's presentation
- 3) Bolus fluid in less than 10 minutes.
- 4) Repeat vital signs and lung auscultation before and after fluid administration.
- 5) If pulsatile abdominal mass present or suspected AAA/TAA, PATCH.

### PEDIATRIC BRADYCARDIA, UNSTABLE




- 1) If airway is managed with BVM for greater than 2 minutes, insert 10-16 Fr. OG tube. Gastric decompression allows adequate pulmonary tidal volumes.
- 2) Special considerations may apply in the presence of severe hypothermia.
- 3) Consider IO use if IV access unavailable.
- 4) Dilute 1:1000 Epinephrine with 3-5 ml of NS flush.
- 5) Limited pediatric data; 15 kg or less pediatric electrodes recommended. For greater than 15 kg use adult electrodes.
- 6) Consider Medical Control input to administer Epinephrine IV continuous infusion at a rate of 0.1 to 1 mcg/kg/min.
- 7) rapid transport is essential in these situations. The above procedures should be performed as the patient is being moved towards the hospital.

**PEDIATRIC PULSELESS ARREST**




- 1) Assess rhythm – quick look, only check pulses if there is an organized rhythm present.
- 2) Evaluate airway, intubate if necessary, limit interruption of CPR as much as possible.
- 3) Once patient is successfully intubated perform continuous asynchronous compression (rate 100/min) with ventilations (rate 8-10/min)
- 4) Pulse checks should be done only if EKG indicates a potentially perfusing rhythm, do not interrupt chest compressions, and be very brief.
- 5) Medications should be administered during CPR as soon as possible after rhythm checks.
- 6) Consider possible causes: Hypovolemia, (volume infusion), hypoxia (ventilation/re-evaluation), acidosis (ventilation/re-evaluation), tension pneumothorax (needle decompression), hypothermia, hypoglycemia, drug overdose, cardiac tamponade (volume infusion), massive AMI, hyperkalemia (consider NaHCO<sub>3</sub>, D50W, massive pulmonary embolism).
- 7) If airway managed with BVM > 2 min. insert 10-16 Fr. OG tube after patient has been intubated.
- 8) If patient remains asystolic or other agonal rhythm after successful intubation, initial medications, no reversible causes are identified, and transport has not been initiated, consider termination of resuscitative efforts by order of a physician. Consider interval since arrest.

**PEDIATRIC TACHYCARDIA WITH PULSES**




1) Probable SVT in pediatrics: History incompatible with presentation, P waves absent or abnormal, HR not variable with activity, abrupt rate changes. Infant rate usually > 220 bpm and children usually > 180 bpm.  
 2) Patients often fit in between borderline and critically unstable situations. In these circumstances, a trial of adenosine may be considered but the medic must be prepared for immediate cardioversion.  
 3) the medic should consult Medical Control and consider reducing the Adenosine dosage in patients who are on Dipyridamole (Persantine) and Carbamazepine ( Tegretol)  
 4. Consider 12 lead EKG  
 5) Or biphasic equivalent  
 6) If probable VT contact Medical Control to administer Amiodarone 5mg/kg, max single dose 150 mg over 20 minutes may repeat two more times to a total of 15 mg/kg/day or Lidocaine 1mg/kg every 5-10 minutes to a total of 3 mg/kg.

**PEDIATRIC – NEONATAL RESUSCITATION**




1) If patient is not vigorous and meconium staining is present deep suction mouth and posterior pharynx then nose. Tracheal suctioning may be necessary before stimulating neonate and proceeding with other resuscitative steps. Vigorous- strong respiratory effort, good muscle tone, heart rate > 100 bpm. Depressed- weak or absent respiratory effort, poor muscle tone/limp, heart rate < 100 bpm.  
 2) Tracheal intubation may be considered at several steps. Tracheal tube should be used for tracheal suctioning.  
 3) Utilize IO or umbilical vein if peripheral IV sites inaccessible.

**PEDIATRIC TRAUMA – MUSCULOSKELETAL INJURY**




1) Patients under 15 years of age.  
2) Contact Medical Control to administer Diazepam 0.1mg/kg for large muscle spasm.

## PEDIATRIC TRAUMA – SPINAL INJURY




**PEDIATRIC ENVENOMATION - ARACHNIDS**




1) Attempts to kill or capture insect or bring to ED are not recommended.  
 2) Contact Medical Control to administer Diazepam for severe pain/muscle spasm.  
 3) Careful observation of respiratory status.

**PEDIATRIC AIRWAY (1)**  
Airway Compromise




1) Medical Control contact is not mandatory, however, the medic is encouraged to discuss the situation with Medical Control if he/she is anticipating a Cricothyroidotomy and the clinical situation is such that there is time for Medical Control contact.  
 2) Verify proper tube placement by visualization of the cords and the tube passing through, bulb tube check/air aspiration, technique > 5 years old, chest wall rise, good breath sounds, absence of gastric sounds, and clinical improvement in patient. Surgical Cricothyroidotomy contraindicated in children < 8 years old.  
 3) OG tube placement if child ventilated with BVM for greater than 2 minutes or obvious gastric distention. Patients with head injuries should only have OG tube insertion, NG tube insertion contraindicated.

## PEDIATRIC AIRWAY OBSTRUCTED



1) Verify proper tube placement by bulb tube check / air aspiration (if patient > 5 years old) or chest wall rise, good breath sounds, absence of gastric sounds, and clinical improvement in patient. Surgical Cricothyroidotomy is contraindicated in patients < 8 years old.


## PEDIATRIC RESPIRATORY – UPPER AIRWAY EMERGENCIES CROUP / EPIGLOTTIS



1) BVM with reservoir with 100% O2 is usually adequate to provide ventilation and oxygenation. If ventilation appears clinically inadequate or transport will be greater than 5 minutes, consider intubation


### PEDIATRIC – SEIZURES OF UNKNOWN ETIOLOGY

Prolonged, Repetitive, or Status Epilepticus




1) Obtain blood sample if utilized by receiving facility.  
 2) Administer 0.5 – 1 Gm/kg of Dextrose. For Neonates to one month old administer D10 2mL/kg. For children one month to one year of age administer D10 5-10mL/kg. For children 1-8 years of age, use D25 2-4 mL/kg. If unable to establish IV, give Glucagon 0.5 mg IM.

## PEDIATRIC ABDOMINAL PAIN, NON TRAUMATIC




## PEDIATRIC HYPOTENSION / SHOCK, NON - TRAUMATIC



- 1) BVM with reservoir with 100% O<sub>2</sub> and cricoid pressure is usually adequate to provide ventilation and oxygenation. If ventilation appears clinically inadequate or transport will be greater than 5 minutes, consider intubation.
- 2) If airway managed with BVM for > 2 minutes, insert 10-16 Fr OG tube. Gastric decompression allows adequate pulmonary tidal volumes.
- 3) Rapid transport is of the utmost importance. Advanced life support procedures should be attempted at the scene, but if unsuccessful with a short period of time, the patient should be transported to the nearest appropriate facility without further delay.
- 4) Repeat assessment and lung auscultation before and after each fluid bolus.
- 5) Assess patient and patient symptoms to suggest cause and treat cause.

**PEDIATRIC POISONING / OVERDOSE (1)**



- 1) Patients who are suspected or known to have ingested substances with a suicidal intent may not refuse transport.
- 2) Administer 0.5 – 1 Gm.kg of Dextrose. For neonates administer D 10 2 mL/kg. For children less than one year of age administer D 10 5 – 10 mL/kg. For children 1-8 years of age, use D 25 2-4 mL/kg. If unable to establish IV, give Glucagon 0.5 mg IM.
- 3) Bring bottles / containers if possible. INSPECT SCENE.
- 4) Consider Medical Control input for Sodium Bicarbonate 1-2 mEq/kg for TCA overdose, Atropine 0.05 mg/kg every 2-4 min. for organophosphate exposure.