

**Versed Use for Adult Violent/Agitated Patient  
(Under Standing Orders)**

The primary purpose of this CQI form is to help improve the documentation related to the standing orders for morphine administration.

For each of the following questions, please provide summary data for your system in the last month.

Agency Name: \_\_\_\_\_

Calendar month for data: \_\_\_\_\_

Total # of patient care reports reviewed: \_\_\_\_\_

**NOTE: A "Yes" answer below includes any chart that has the item documented or includes a narrative explanation for why documentation is not provided. A "No" answer includes charts where the item is not documented and not explained.**

	# YES	# NO
1. Vital Signs Documented before and after each administration of Versed?		
2. Documentation of attempts to de-escalate Pt's behavior prior to restraining?		
3. Documentation of type of restraint used?		
4. Documentation of appropriate position of restraint?		
5. Documentation of distal CMS q 10 minutes for extremities restrained?		
3. Correct dose of <b>IM or IN</b> Versed? (5mg <b>IM or IN</b> age < 60 and 2.5mg IM or IN in age > 60)		
4. Dose repeated x 1 in 10 minutes if inadequate response to initial dose? ( <i>should not be repeated sooner than 10 minutes</i> )		
5. If patient remains combative or violent and <b>IV access</b> is obtained, correct dose of <b>IV</b> Versed given? (Versed 2.5mg IV slow push over 2 min. for age <60 and Versed 1-3mg IV slow push over 2 min if age > 60, may repeat x 1 in 10 min.)		
6. 1 liter bolus given if IV access obtained?		
7. Patch made for authorization of soft restraints?		

Criteria Met \_\_\_\_\_ Criteria Not Met \_\_\_\_\_

# of patients that required assisted ventilations after any administration? \_\_\_\_\_

# of patients that required Narcan administration? \_\_\_\_\_

# of patients that had hypotension: \_\_\_\_\_